

Recovery Strategy and Action Plan for the Fawnsfoot (*Truncilla donaciformis*) and Threehorn Wartyback (*Obliquaria reflexa*) in Canada

Fawnsfoot



Threehorn Wartyback



2020

Recommended citation:

Fisheries and Oceans Canada. 2020. Recovery Strategy and Action Plan for the Fawnsfoot (*Truncilla donaciformis*) and Threehorn Wartyback (*Obliquaria reflexa*) in Canada [Proposed]. *Species at Risk Act* Recovery Strategy Series. Fisheries and Oceans Canada, Ottawa. vi + 66 pp.

For copies of the recovery strategy and action plan, or for additional information on species at risk, including Committee on the Status of Endangered Wildlife in Canada (COSEWIC) status reports, residence descriptions, action plans, and other related recovery documents, please visit the [Species at Risk Public Registry](#).

Cover illustration: Photo courtesy of the National Water Research Institute

Également disponible en français sous le titre

« Programme de rétablissement et plan d'action pour la troncille pied-de-faon (*Truncilla donaciformis*) et l'obliquaire à trois cornes (*Obliquaria reflexa*) au Canada »

© Her Majesty the Queen in Right of Canada, represented by the Minister of Fisheries and Oceans, 2020. All rights reserved.

ISBN ISBN to come

Catalogue no. Catalogue no. to come

Content (excluding the illustrations) may be used without permission, with appropriate credit to the source.

Preface

The federal, provincial, and territorial government signatories under the [Accord for the Protection of Species at Risk \(1996\)](#) agreed to establish complementary legislation and programs that provide for effective protection of species at risk throughout Canada. Under the *Species at Risk Act* (S.C. 2002, c.29) (SARA), the federal competent ministers are responsible for the preparation of a recovery strategy and action plan for species listed as extirpated, endangered, or threatened and are required to report on progress five years after the publication of the final document on the Species at Risk Public Registry.

This document has been prepared to meet the requirements under SARA of both a recovery strategy and an action plan. As such, it provides both the strategic direction for the recovery of the species, including the population and distribution objectives for the species, as well as the more detailed recovery measures to support this strategic direction, outlining what needs to be done to achieve the objectives. SARA requires that an action plan also include an evaluation of the socio-economic costs that may be incurred by the more detailed recovery measures, as well as the benefits to be derived from its implementation. It is important to note that the setting of population and distribution objectives and the identification of critical habitat are science-based exercises, therefore, socio-economic factors were not considered in their development. The socio-economic evaluation only applies to the more detailed recovery measures (that is, the action plan portion).

The Minister of Fisheries and Oceans is the competent minister under SARA for the Fawnsfoot and Threehorn Wartyback and has prepared this recovery strategy and action plan, as per sections 37 and 47 of SARA. In preparing this recovery strategy and action plan, the competent minister has considered, as per section 38 of SARA, the commitment of the Government of Canada to conserving biological diversity and to the principle that, if there are threats of serious or irreversible damage to the listed species, cost-effective measures to prevent the reduction or loss of the species should not be postponed for a lack of full scientific certainty. To the extent possible, this recovery strategy and action plan has been prepared in cooperation with the Ontario Ministry of Natural Resources and Forestry, Environment and Climate Change Canada, and conservation authorities who manage watersheds where these species are present, as per section 39(1) and 48(1) of SARA.

As stated in the preamble to SARA, success in the recovery of these species depends on the commitment and cooperation of many different constituencies that will be involved in implementing the directions set out in this recovery strategy and action plan and will not be achieved by Fisheries and Oceans Canada, or any other jurisdiction alone. The cost of conserving species at risk is shared amongst different constituencies. All Canadians are invited to join in supporting and implementing this recovery strategy and action plan for the benefit of the Fawnsfoot and Threehorn Wartyback and Canadian society as a whole.

Acknowledgments

Fisheries and Oceans Canada (DFO) would like to thank the contributing authors, Kelly McNichols-O'Rourke (DFO), Pat Dimond (DFO contractor), Peter Jarvis (DFO contractor), Joshua Stacey (DFO), and Amy Boyko (DFO), as well as the following organizations for their support in the development of the Fawnsfoot and Threehorn Wartyback recovery strategy and action plan: Ontario Freshwater Mussel Recovery Team, Environment and Climate Change Canada, Ontario Ministry of Natural Resources and Forestry, Upper Thames River

Conservation Authority, Lower Thames Valley Conservation Authority, and the Bishop Mills Natural History Centre.

Executive summary

The Fawnsfoot and Threehorn Wartyback were listed as Endangered and Threatened, respectively, under the *Species at Risk Act* (SARA) in August 2019. This multispecies recovery strategy and action plan is considered one in a series of documents for these species that are linked and should be taken into consideration together, including: the Committee on the Status of Endangered Wildlife in Canada (COSEWIC) status reports for Fawnsfoot (2008) and Threehorn Wartyback (2013); and, the recovery potential assessments for the Fawnsfoot (2011) and Threehorn Wartyback (2014). Recovery for these species has been determined to be biologically and technically feasible.

The Fawnsfoot is a small freshwater mussel of approximately 35 mm in length with a moderately thick, oval to triangular shell that is rounded on the anterior end and bluntly pointed on the posterior. The shell is smooth, yellow to greenish in colour and has dark green rays that are often broken into v-shaped or chevron markings. The Threehorn Wartyback is a medium-sized freshwater mussel restricted to central North America from the Gulf of Mexico to the Great Lakes watershed. Its thick shell can be green, tan or brown with a maximum length of 80 mm, and is circular to triangular in shape, while the anterior end is rounded and the posterior end is bluntly pointed. The defining characteristic of this species is the single row of two to five large knobs or “horns”, which distinguishes this species from other freshwater mussels found in Canada.

While freshwater mussels are among the world’s most imperilled taxa, southern Ontario remains home to the largest and most diverse freshwater mussel communities in Canada. The Canadian distribution of both of these species is restricted to southern Ontario in the Great Lakes watershed. Both the Fawnsfoot and Threehorn Wartyback are currently present in the Grand, Sydenham, and Thames rivers, while the Fawnsfoot is also distributed within the delta area of the St. Clair River, and Threehorn Wartyback has recently been captured in the Detroit River. Overall, the Canadian ranges of both species have diminished in comparison with their historical distributions as they are no longer encountered in Lake Erie, and in the case of the Threehorn Wartyback, Lake St. Clair. However, the Threehorn Wartyback appears to have never been a major component of the mussel fauna in Canada.

The main threats facing the species include: the presence of invasive species (primarily dreissenid mussels [Zebra Mussel and Quagga Mussel]); turbidity, sediment and nutrient loading; contaminants and toxic substances; habitat removal/alteration; altered flow regimes; predation and harvesting; declines in host fish availability; and, recreational activities (for example, ATVs, boat propellers, paddling).

The population and distribution objectives for Fawnsfoot are to return self-sustaining populations in the lower Grand River, the St. Clair River delta, the East Sydenham River, the North Sydenham River (Bear Creek), and the lower Thames River. The population and distribution objectives for Threehorn Wartyback are to return self-sustaining populations in the lower Grand River, the East Sydenham River, the North Sydenham River (Bear Creek) and the lower Thames River. The populations at these locations could be considered recovered when they demonstrate active signs of reproduction and recruitment throughout their distribution and are stable or increasing with low risk from known threats.

A description of the broad strategies to be taken to address threats to the species' survival and recovery, as well as research and management approaches needed to meet the population and distribution objectives are included in section 7.

Using the best available information, critical habitat has been identified for these species to the extent possible, and provides the functions and features necessary to support their life-cycle processes and to achieve their population and distribution objectives. This recovery strategy and action plan currently identifies critical habitat for Fawnsfoot and Threehorn Wartyback in the North Sydenham (Bear Creek), East Sydenham, Grand, and Thames rivers. A schedule of studies has been developed that outlines the necessary steps to obtain the required information to further refine these critical habitat descriptions. It is anticipated that the protection of these species' critical habitat from destruction will be accomplished through a SARA critical habitat order made under subsections 58(4) and (5), which will invoke the prohibition in subsection 58(1) against the destruction of the identified critical habitat.

The action plan portion of this document provides the detailed recovery planning in support of the strategic direction set out in the recovery strategy section of the document. The action plan outlines what needs to be done to achieve the population and distribution objectives, including the measures to be taken to address threats and monitor the recovery of the species, as well as the required measures to protect critical habitat. Socio-economic impacts of implementing the action plan are also evaluated.

Recovery feasibility summary

Recovery of the Fawnsfoot and Threehorn Wartyback is believed to be biologically and technically feasible. Recovery feasibility is determined according to four criteria by the Government of Canada (2009):

1. Are individuals of the wildlife species that are capable of reproduction currently available now or in the foreseeable future to sustain the population or improve its abundance?

Yes. Reproducing populations of both Fawnsfoot and Threehorn Wartyback exist in at least the East Sydenham and Thames rivers. These are available to improve the population growth rate and abundance.

2. Is sufficient suitable habitat available to support these species or could it be made available through habitat management or restoration?

Yes. The habitat that supports these species is sufficient but in some locations it is of marginal quality due to the presence of dreissenid mussels. At locations with declining populations, suitable habitat may be made available through current and proposed restoration efforts. With respect to Fawnsfoot, the St. Clair River delta population receives some protection from human disturbance as it is located within Walpole Island First Nation reserve where permits are required to access these areas.

3. Can significant threats to the species or its habitat be avoided or mitigated?

Yes. With the exception of dreissenid mussels in the Great Lakes, significant threats to populations of both species, such as sedimentation and nutrient and contaminant loading, can be avoided or mitigated through recovery actions including many activities that are already underway. While action has been taken to limit the expansion of dreissenid mussels, recovery in heavily infested areas (for example, Lake St. Clair, Detroit River and Lake Erie) to historical levels is not possible; however, currently existing refuge sites at these locations should be maintained.

4. Do recovery techniques exist to achieve the population and distribution objectives or can they be developed within a reasonable timeframe?

Yes. Recovery techniques that are necessary to recover Fawnsfoot and Threehorn Wartyback populations do exist and have been demonstrated to be effective. For example, artificial propagation in the U.S. has been successful for a number of species (Hanlon 2000), while similarly, it may be possible to artificially propagate juveniles of the host fish species that have been identified. In addition, techniques for the reduction of identified threats (for example, best management practices to reduce sedimentation) and restoration of habitats are also well documented as effective recovery measures. For example, actions to improve water quality and fish movement (important for host fish populations) have resulted in an increase in the species richness of freshwater mussels in the Grand River (Metcalfe-Smith et al. 2000a). It is important to note that the effort expended to achieve recovery will not be uniform across all locations; much greater effort will be required to improve habitat in areas with reduced populations.

Table of contents

Preface.....	i
Acknowledgments	i
Executive summary	iii
Recovery feasibility summary.....	v
Background	1
1. Introduction	1
2. COSEWIC species assessment information	2
3. Species status information	3
4. Species information.....	4
4.1 Description.....	4
4.2 Population abundance and distribution.....	6
4.3 Needs of the species	13
5. Threats.....	14
5.1 Threat assessment	14
5.2 Description of threats.....	15
Recovery	20
6. Population and distribution objectives	20
7. Broad strategies and general approaches to meet objectives.....	21
7.1 Actions already completed or currently underway	21
7.2 Measures to be taken to implement the recovery strategy/action plan	22
7.3 Narrative to support the recovery planning and implementation tables	30
8. Critical habitat	33
8.1 Identification of Fawnsfoot and Threehorn Wartyback critical habitat	33
8.1.1. General description of Fawnsfoot and Threehorn Wartyback critical habitat .	33
.....	33
8.1.2. Information and methods used to identify critical habitat	34
8.1.3. Identification of critical habitat.....	34
8.2 Schedule of studies to identify critical habitat	48
8.3 Activities likely to result in the destruction of critical habitat.....	48
8.4 Proposed measures to protect critical habitat.....	54
9. Evaluation of the socio-economic costs and of benefits of the action plan.....	54
10. Measuring progress	56
11. References.....	57
Appendix A: effects on the environment and other species.....	65
Appendix B: record of cooperation and consultation	66

Background

1. Introduction

The Fawnsfoot (*Truncilla donaciformis*) was listed as Endangered under the *Species at Risk Act* (SARA) in August 2019. The Threehorn Wartyback (*Obliquaria reflexa*) was listed as Threatened under SARA in August 2019. This recovery strategy and action plan is part of a series of documents regarding the Fawnsfoot and Threehorn Wartyback that should be taken into consideration together, including the Committee on the Status of Endangered Wildlife in Canada (COSEWIC) status reports for Fawnsfoot ([COSEWIC 2008](#)) and Threehorn Wartyback ([COSEWIC 2013](#)), and the science advisory reports from the recovery potential assessments (RPA) for Fawnsfoot ([DFO 2011](#)), and Threehorn Wartyback ([DFO 2014](#)).

A recovery strategy is a planning document that identifies what needs to be done to arrest or reverse the decline of a species in addition to providing background information on the species and its threats and critical habitat information. It sets objectives and identifies the main areas of activities to be undertaken, while the action plan portion provides the detailed recovery planning that supports the strategic direction set out in the recovery strategy portion. Action planning for species at risk recovery is an iterative process; therefore, the implementation schedule (tables 7 to 9) in this recovery strategy and action plan may be modified in the future depending on the progression towards recovery.

The RPA is a process developed by Fisheries and Oceans Canada (DFO) Science to provide the information and scientific advice required to implement SARA, relying on the best available scientific information, data analyses and modelling, and expert opinions. The outcome of this process informs many sections of the recovery strategy and action plan. For more detailed information beyond what is presented in the recovery strategy and action plan, refer to the COSEWIC status reports and the RPA science advisory reports.

2. COSEWIC species assessment information

Date of assessment: April 2008

Species' common name (population): Fawnsfoot

Scientific name: *Truncilla donaciformis*

Status: Endangered

Reason(s) for designation: This freshwater mussel is widely distributed in central North America, with the northern portion of its range extending into the Lake Erie, Lake St. Clair and lower Lake Huron drainages of southwestern Ontario. It appears to have always been a rare species in Canada, representing < 5% of the freshwater mussel community in terms of abundance wherever it occurs. Approximately 86% of historical records are in waters that are now infested with zebra mussels and therefore uninhabitable. Zebra mussels, which were accidentally introduced into the Great Lakes, attach to the shells of native freshwater mussels, causing them to suffocate or die from lack of food. The species has declined dramatically and has been lost from four historical locations resulting in a 51% reduction in its range. It is now found in only five widely separated locations, two of which represent single specimens. In two locations, the species' distribution may be limited by the presence of dams that restrict the movements of the freshwater drum, the presumed fish host of the juvenile mussels. Poor water quality resulting from rural and urban influences poses an additional continuing threat.

Canadian occurrence: Ontario

Status history: Designated Endangered in April 2008. Assessment based on a new report.

Date of assessment: May 2013

Species' common name (population): Threehorn Wartyback

Scientific name: *Obliquaria reflexa*

Status: Threatened

Reason(s) for designation: This rare species historically occurred in the Great Lakes drainages including Lake St. Clair, western Lake Erie, and the Grand, Thames, and Detroit rivers. The species has not been found since 1992 in Lake St. Clair and the Detroit River and may be extirpated there due largely to the impacts of Zebra and Quagga mussels. It was last recorded from the Canadian side of Lake Erie in 1997. Pollution (sediment loading, nutrient loading, contaminants and toxic substances) related to both urban and agricultural activities represents a high and continuing threat at the three remaining riverine locations.

Canadian occurrence: Ontario

Status history: Designated Threatened in May 2013.

3. Species status information

Table 1. Summary of existing protection or other status designations assigned to Fawnsfoot

Jurisdiction	Authority/organization	Year(s) assessed and/or listed	Status/description	Designation level
Ontario	<i>Endangered Species Act 2007*</i>	2009	Endangered	Population
Ontario	NatureServe	2015	Provincial: S2, Imperilled	Population
Canada	Committee on the Status of Endangered Wildlife in Canada (COSEWIC)	2008	Endangered	Population
Canada	<i>Species at Risk Act (SARA)</i>	2019	Endangered	Population
Canada	NatureServe	2010	National: N2, Imperilled	Population
United States ¹	NatureServe	1998	National: N5, Secure	Population
International	NatureServe	2011	Global: G5, Secure	Species

Table 2. Summary of existing protection or other status designations assigned to Threehorn Wartyback

Jurisdiction	Authority/organization	Year(s) assessed and/or listed	Status/description	Designation level
Ontario	<i>Ontario's Endangered Species Act 2007*</i>	2014	Threatened	Population
Ontario	NatureServe	2013	Provincial: S1, Critically Imperilled	Species
Canada	Committee on the Status of Endangered Wildlife in Canada (COSEWIC)	2013	Threatened	Species
Canada	<i>Species at Risk Act (SARA)</i>	2019	Threatened	Species

¹ Refer to NatureServe 2016 for state specific designations

Jurisdiction	Authority/organization	Year(s) assessed and/or listed	Status/description	Designation level
Canada	NatureServe	2013	National: N1, Critically Imperilled	Population
United States	NatureServe	1998	National: N5, Secure	Population
International	NatureServe	2007	Global: G5, Secure	Species

*Under the Act, individuals are currently protected and their habitat has been protected under the general habitat protection provisions of the Act since 2009 and 2014 for Fawnsfoot and Threehorn Wartyback, respectively.

The listing of Fawnsfoot and Threehorn Wartyback as Endangered and Threatened species (respectively) provides immediate protection wherever these species are found in Canada by section 32 of SARA:

“No person shall kill, harm, harass, capture or take an individual of a wildlife species that is listed as an extirpated species, an endangered species or a threatened species.”
[subsection 32(1)]

“No person shall possess, collect, buy, sell or trade an individual of a wildlife species that is listed as an extirpated species, an endangered species or a threatened species, or any part or derivative of such an individual.” [subsection 32(2)]

Under section 73 of SARA, the competent minister may enter into an agreement or issue a permit authorizing a person to engage in an activity affecting a listed wildlife species, any part of its critical habitat or its residences.

4. Species information

4.1 Description

Fawnsfoot

The Fawnsfoot is a small freshwater mussel approximately 35 mm in length (COSEWIC 2008). The shell is moderately thick, oval to triangular, rounded on the anterior end and bluntly pointed on the posterior (COSEWIC 2008). The shell is smooth, yellow to greenish in colour and has dark green rays that are often broken into v-shaped or chevron markings (COSEWIC 2008). The beaks (the oldest part of the shell) are full, central, and slightly elevated above the hinge line and have 3 to 8 fine bars (COSEWIC 2008). Fawnsfoot is generally found in deeper areas (1 to >5 m) in large, slow- to moderate-flowing rivers, although it may also inhabit lakes and reservoirs (COSEWIC 2008).



Figure 1. Fawnsfoot. Photograph by Environment and Climate Change Canada, reproduced with permission.

Threehorn Wartyback

As described by Watters et al. (2009), Metcalfe-Smith et al. (2005), Clarke (1981) and COSEWIC (2013), the Threehorn Wartyback's shell is generally green, tan or brown with a maximum length of 80 mm, although lengths to 40 mm are more common. The thick shell is circular to triangular in shape, while the anterior end is rounded and the posterior end is bluntly pointed. The defining characteristic of this species is the single row of two to five large knobs or "horns", which distinguishes this species from other freshwater mussels found in Canada. The Threehorn Wartyback is the only member of the genus *Obliquaria* that occurs in Canada.




Figure 2. Threehorn Wartyback. Photograph by Environment and Climate Change Canada, reproduced with permission.

Ecological role: Both the Fawnsfoot and Threehorn Wartyback, like other freshwater mussels, play an integral role in the functioning of aquatic ecosystems, including water column and sediment processes (Vaughn and Hakenkamp 2001) as well as food web dynamics, linking and influencing multiple trophic levels (for example, Vaughn et al. 2004; Vaughn and Spooner 2006). Furthermore, their filtration of suspended materials facilitates the transfer of energy and nutrients from the water column to the sediment. Mussels are sensitive indicators of the health of freshwater ecosystems, including water and habitat quality and, especially, the fish community on which they depend for successful reproduction. Mussels can provide habitat for other organisms by providing physical structure, and dense mussel beds can stabilize streambed substrates during periods of high flow. Rare species, including other unionid mussels, have been shown to benefit energetically from living in species-rich communities (Spooner 2007). Freshwater mussels are also important prey for several species including the Muskrat (*Ondatra zibethicus*) (Neves and Odom 1989), which results in a transfer of energy from the aquatic to the terrestrial environment. More detailed information can be found in COSEWIC (2008) and COSEWIC (2013).

4.2 Population abundance and distribution

Global distribution and population abundance

Fawnsfoot: In the U.S., Fawnsfoot is considered secure and occurs in central North America (Alabama, Arkansas, Georgia, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Michigan, Minnesota, Mississippi, Missouri, Nebraska, New York, Ohio, Oklahoma, Pennsylvania, South Dakota, Tennessee, Texas, West Virginia, and Wisconsin) (figure 3). The current distribution of Fawnsfoot is similar to its historical distribution, but the species has declined in many places, particularly in Lake Erie (NatureServe 2015). In Canada, Fawnsfoot occurs only in Ontario (figure 4).

Threehorn Wartyback: The Threehorn Wartyback occurs throughout most of the Mississippi River drainage as well as the state of Michigan and province of Ontario (figures 4 and 6). The species' range encapsulates southwestern Ontario, western Pennsylvania, Minnesota, eastern Kansas, Oklahoma, Texas, and the Coosa-Alabama River and Tombigbee River systems in the southeastern U.S. (figure 4). Although considered stable throughout its global range, Threehorn Wartyback appears to be extirpated from the offshore waters of Lake St. Clair and the Canadian side of Lake Erie (NatureServe 2015).

Canadian distribution and population abundance

Fawnsfoot: In Canada, Fawnsfoot is known only from the Great Lakes watershed of Ontario. The current distribution of the species includes the Grand, Sydenham, and Thames rivers, Muskrat Creek (of the Saugeen River), the St. Clair River delta (COSEWIC 2008), and potentially the Welland River (figure 5). The range of Fawnsfoot has been significantly reduced by ~51% in Canada as it is believed to have been extirpated from the Detroit and Niagara rivers, Lake Erie and the offshore waters of Lake St. Clair. For more information refer to the COSEWIC status report and the RPA science advisory report.

Threehorn Wartyback: In Canada, Threehorn Wartyback, like Fawnsfoot, is found in the Grand, Sydenham, and Thames rivers within the Great Lakes watershed of southern Ontario (figure 6). As with Fawnsfoot, the distribution of Threehorn Wartyback is believed to have undergone a contraction due to the establishment of invasive dreissenid mussels (Zebra Mussel

[*Dreissena polymorpha*] and Quagga Mussel [*Dreissena bugensis*] in historical locations (that is, Lake St. Clair, and western Lake Erie) (COSEWIC 2013).




Figure 3. Global distribution (shaded area) of the Fawnsfoot (COSEWIC 2008)




Figure 4. Global distribution of the Threehorn Wartyback (from COSEWIC 2013)




Figure 5. Historical (pre-1996) and current (1996 to 2019) distribution of Fawnsfoot in Canada




Figure 6. Historical (pre-1996) and current (1996 to 2019) distribution of Threehorn Wartyback in Canada

Population assessment

Fawnsfoot: To date, it appears that there are five remaining populations of Fawnsfoot in Ontario. The largest population occurs in the lower portion of the Thames River, while others can be found in the Grand and East and North Sydenham rivers. In addition, very small populations are thought to exist in the St. Clair River delta and were thought to exist in and Muskrat Creek (a tributary of the Teeswater River in the Saugeen River watershed) as only one individual has ever been found in both systems. However, after recent sampling in 2019 it seems unlikely that Fawnsfoot still occurs in Muskrat Creek and there is uncertainty as to whether a population ever existed in this waterbody or alternatively the specimen detected had been inadvertently transferred to this location. The species may also be present in the Welland River and its Feeder Canal; however, further research is pending to confirm occupancy at these two locations.

Historically, Fawnsfoot was recorded in the Great Lakes watershed including the Detroit River and Lake St. Clair, the St. Clair River delta, and the Grand, Sydenham, and Thames rivers. For more information regarding Fawnsfoot records within these waters, refer to the RPA science advisory report (DFO 2011). More recently, Fawnsfoot has been discovered in the Saugeen River, North Sydenham River (Bear Creek) and potentially the Welland River. Since the drafting of the RPA, which was published in 2011, updated information regarding the distribution and occurrence of Fawnsfoot has become available as a result of further detections. For example, within the 2011 to 2016 time period, four live specimens have been detected in the Grand River, with one detection occurring just downstream of the Caledonia Dam; 12 live specimens have been detected in the East Sydenham River; and, 201 live individuals were recorded during surveys in the Thames River. Furthermore, one weathered shell and one live specimen, of which it is unclear if it was a Fawnsfoot or Deertoe (*Truncilla truncata*), were detected in the Welland River in 2015 and 2016, respectively. One weathered shell was also discovered in the Feeder Canal, which historically connected the Grand and Welland river watersheds; however, the source of this shell (potential human transport) is uncertain. In addition, a live specimen was captured in Bear Creek, a tributary of the North Sydenham River, for the first time in 2016. Although no live specimens were detected, six weathered shells were found in Rondeau Bay.

Overall, Fawnsfoot is believed to be declining (COSEWIC 2008; NatureServe 2015) throughout its historical range. Although Fawnsfoot has always been a small component of the mussel community (< 5% wherever it occurs), it has declined by 51% in its extent of occurrence primarily due to the establishment of dreissenid mussels among other threats.

Populations of Fawnsfoot were ranked by Bouvier and Morris (2011), with respect to their abundance and trajectory. Population abundance and trajectories were then combined to determine the population status (table 3). A certainty level was also assigned to the population status, which reflected the lowest level of certainty associated with either population abundance or trajectory. Fawnsfoot populations are either extirpated or in poor health (declining) with the exception of the Thames River population, which is considered to be fair in status. Refer to Bouvier and Morris (2011) for further results and details regarding the methodology. The North Sydenham River (Bear Creek) populations were not known at the time that the RPA was being developed; therefore this location is not included in table 3. The status and trajectory of Fawnsfoot populations at this location is currently unknown.

Table 3. Population status of Fawnsfoot populations in Canada, resulting from an analysis of both the Relative Abundance Index and Population Trajectory (table modified from Bouvier and Morris 2011)

Population	Population status	Certainty*
Grand River	Poor	3
Great Lakes and connecting channels	Extirpated	2
Saugeen River**	Poor	3
St. Clair River delta**	Poor	3
Sydenham River	Poor	3
Thames River	Fair	3

* Certainty assigned to each population status reflects the lowest level of certainty associated with either abundance index or population trajectory. Certainty associated with abundance index or population trajectory is listed as: 1=quantitative analysis; 2=standardized sampling; 3=expert opinion.

** Population represented by a single live individual.

Threehorn Wartyback: Threehorn Wartyback has always been a rare species in the Canadian faunal record (COSEWIC 2013). At present, it occurs in the Grand, East Sydenham, North Sydenham (Bear Creek) and Thames rivers, with the most frequent detections over the last 28 years occurring in the Sydenham River (Bouvier et al. 2014). For further information regarding Threehorn Wartyback records within these waters, refer to the RPA science advisory report (DFO 2014).

As discussed for Fawnsfoot, a contraction of the distribution of Threehorn Wartyback is thought to have occurred based on a lack of survey detections in historical locations (that is, Lake St. Clair, western Lake Erie, and Detroit River). This species is considered extirpated from the Canadian side of Lake Erie; although, one fresh shell was reported in 2001, and four weathered valves were discovered in 2014 from Rondeau Bay (Lower Great Lakes Unionid Database 2016). A total of 34 live specimens were captured from two sites in the Detroit River in 2019 between the mouth of the Canard River and Edgewater Beach representing the first detections in this waterbody in decades.

Within the East Sydenham River, one site located at Dawn Mills has been re-sampled annually from 2002 to 2009 and has resulted in the observance of 72 live individuals (33 recaptures and 39 new individuals). From 2010 to 2018, a total of 13 live specimens have been captured in the East Sydenham River. It is believed that recruitment is occurring in the Sydenham River population based on the current size frequency distribution, and the observation of a 15 mm individual (K. McNichols-O'Rourke, DFO, pers. obs.). The species was also detected in the North Sydenham River (Bear Creek) for the first time in 2018 when eight live individuals were discovered by SCRCA. Threehorn Wartyback is currently known to occupy a 100 km reach of the Thames River and a total of 30 live individuals have been collected from this system in the 1998 to 2010 time period (refer to RPA science advisory report DFO 2013). Since that time 20 live specimens have been captured. Within the Grand River, four live individuals, five fresh shells and seven weathered shells were recorded from seven sites sampled in 2011 (refer to RPA science advisory report DFO 2013). Since 2011, eight live individuals were detected in Mazi Drain, four live individuals were detected downstream of Dunnville, and three live individuals were captured downstream of Cayuga.

Populations of Threehorn Wartyback have also been ranked in the same manner described above for Fawnsfoot (table 4). Refer to DFO (2014) for details on the methods used in the assessment of Population Status. Based on these results, the status of Threehorn Wartyback populations in the Grand, Sydenham, and Thames rivers is believed to be poor; however, there is evidence of recruitment in the Sydenham and Thames rivers based on size frequency distributions of recent collections. Unfortunately, too few live specimens have been encountered in the Grand River to comment on recruitment at this time. As mentioned for Fawnsfoot, the North Sydenham River is not included in this table, therefore the status and trajectory of this population is currently unknown.

Table 4. Population Status of all Threehorn Wartyback populations in Canada, resulting from an analysis of both the Relative Abundance Index and Population Trajectory (table adapted from DFO 2014)

Population	Population status	Certainty*
Sydenham River	Poor	3
Thames River	Poor	3
Detroit River	Unknown	N/A
Grand River	Poor	3
Great Lakes	Extirpated	N/A

* Certainty assigned to each population status is reflective of the lowest level of certainty associated with either the abundance index or population trajectory. Certainty associated with abundance index or population trajectory is listed as: 1=quantitative analysis; 2=standardized sampling; 3=expert opinion.

4.3 Needs of the species

Habitat and biological needs

As studies specific to the biological requirements of the Fawnsfoot and Threehorn Wartyback are rare, conclusions must be drawn from generalized habitat requirements of unionids to define their needs. Both species, like other freshwater mussels belonging to the Unionidae family, exhibit complex life cycles that are dependent on both environmental and biological components. For example, larvae, known as glochidia, are released from the female's gills and uptaken by a suitable fish species (host fish) where they become encysted on the fishes' gills and feed on body fluids until they metamorphose into juveniles (COSEWIC 2008). After metamorphosis, juveniles release themselves from the host and fall to the substrate to begin life as free-living mussels. Juvenile mussels remain buried until they are sexually mature, at which point they move to the surface for the dispersal/intake of gametes (Watters et al. 2001).

Fawnsfoot: The most likely host for Fawnsfoot in Canada is Freshwater Drum (*Aplodinotus grunniens*), although Sauger (*Sander canadensis*) has also been reported as a potential host (Surber 1913; Wilson 1916; Clarke 1981). Adult Fawnsfoot are usually found in substrates of sand or mud (Clark 1981; Parmalee and Bogan 1998), but can be found in areas with coarser substrates (Howells et al. 1996). Remaining populations in Canada are usually found in the lower portions of larger rivers on fine sand or gravel substrates.

Threehorn Wartyback: Although the host fish(es) have not been identified for Canadian populations, four fish species have been identified as hosts in U.S. populations, including Common Shiner (*Luxilus cornutus*), Longnose Dace (*Rhinichthys cataractae*), Goldeye (*Hiodon alosoides*), and Silverjaw Minnow (*Notropis buccatus*) (Barnhart and Baird 2000; Watters et al. 2009). All but Silverjaw Minnow have been found in Ontario, with Common Shiner and

Longnose Dace confirmed to overlap Threehorn Wartyback's Canadian distribution (Holm et al. 2009). The Threehorn Wartyback appears most commonly in large rivers with moderate current and in shallow embayments and reservoirs with little current (Clarke 1981; Metcalfe-Smith et al. 2005; Watters et al. 2009). The species has been observed in a variety of substrate types (that is, clay, detritus, silt, sand, gravel, rubble and boulder); although, sand and gravel seem to be preferred (see COSEWIC 2013).

It is clear that the needs of both the Fawnsfoot and the Threehorn Wartyback are similar in terms of reproduction and they share a general preference for sand, mud and gravel substrates. Taken together, this information indicates that both species require available host fish(es) and pervious substrates that allow juveniles to burrow and adults to embed themselves. For further information on the habitat and biological needs of the Fawnsfoot and Threehorn Wartyback, refer to the COSEWIC status reports and the RPA science advisory reports for these species.

Limiting factors for both species:

- predation
- reliance on host fishes
- largely sedentary existence for juvenile and adult stages, hence limited ability to disperse and to relocate from substandard conditions

5. Threats

5.1 Threat assessment

Table 5 adapted from the Fawnsfoot RPA (DFO 2011), and table 6 from the Threehorn Wartyback RPA (DFO 2014), provide summaries of the threats to Fawnsfoot and Threehorn Wartyback populations in Canada, respectively. Known and suspected threats were ranked with respect to threat likelihood and impact for each population. The threat likelihood and impact categories were then combined to produce an overall threat level. A certainty level was also assigned to the overall threat level, which reflected the lowest level of certainty associated with either threat likelihood or threat impact. See DFO (2011) and DFO (2014) for further details.

Table 5. Threat levels for Canadian Fawnsfoot populations

Threat ²	Sydenham River	Lower Thames River	Grand River	St. Clair River delta	Saugeen River
Invasive species	Medium (2)	High (2)	High (2)	High (2)	Medium (2)
Turbidity and sediment loading	Medium (3)	Medium (3)	Medium (2)	Low (3)	High (3)
Contaminants and toxic substances	High (3)	High (3)	High (2)	High (3)	High (3)
Nutrient loading	Medium (3)	Medium (3)	Medium (2)	Low (3)	High (3)
Altered flow regimes	Medium (3)	Medium (3)	Medium (2)	Unknown	Medium (3)

² N.B. The threat level represents a combination of the current threat impact and threat likelihood at a location. It does not reflect the potential impact a threat might have on a freshwater mussel population if it was allowed to occur in the future.

Threat ²	Sydenham River	Lower Thames River	Grand River	St. Clair River delta	Saugeen River
Habitat removal and alterations	High (3)	High (3)	High (2)	Medium (3)	High (3)
Host fish(es)	Medium (3)	Medium (3)	High (3)	Medium (3)	Unknown
Recreational activities	Low (3)	Low (3)	Low (3)	Low (3)	Low (3)

Table 6. Threat level for Canadian Threehorn Wartyback populations

Threats	Sydenham River	Thames River	Grand River
Invasive species	Low (2)	High (2)	High (2)
Turbidity	Medium (3)	Unknown (3)	Unknown (3)
Sediment loading	Medium (3)	Medium (3)	Medium (3)
Contaminants and toxic substances	High (3)	High (3)	High (3)
Nutrient loading	Medium (3)	Medium (3)	High (3)
Altered flow regimes	Low (3)	Low (3)	Medium (3)
Habitat removal and alteration	High (3)	High (3)	High (3)
Host fish decline (due to invasive species)	Unknown (3)	Unknown (3)	Unknown (3)

5.2 Description of threats

Invasive species: Dreissenid mussels have decimated populations of freshwater mussels, including Fawnsfoot and Threehorn Wartyback, in the lower Great Lakes by virtually eliminating historical habitat (Gillis and Mackie 1994; Schloesser and Nalepa 1994; Nalepa et al. 1996). For example, approximately 86% of historical records for Fawnsfoot and Threehorn Wartyback are from areas now infested with dreissenid mussels, making them uninhabitable. In addition, further expansion of dreissenids in the St. Clair River delta continues to threaten and limit the distribution of both these species.

Dreissenid mussels have been observed to colonize the shells of native unionids in large numbers, inhibiting their ability to open and close valves; limiting their movement, feeding, burrowing and reproductive activities; and, increasing their risk of predation and parasitism (Schloesser et al. 1996; Baker and Hornbach 1997). Additionally, dreissenid mussels have been shown to directly reduce available food sources in the water column, which can directly impact the fitness of unionids due to the similarity of their diets (Mackie 1991). The natural dispersal of dreissenid mussels is passive and generally occurs downstream of the adult population during the larval stage, via water currents in lentic environments; however, it can also threaten riverine mussel populations if it is introduced to upstream locations such as reservoirs (Bouvier and Morris 2011).

Currently, dreissenid mussels have been reported in the Springbank and Fanshawe reservoirs on the Thames River (UTRCA 2012) and throughout the lower Thames River (Morris and Edwards 2007). Although dreissenids have only been detected in the Grand River south of the Dunnville Dam (S. Staton, DFO, pers. comm. 2015), the freshwater mussel populations therein

are highly susceptible to invasion, as the lower river is heavily impounded. Furthermore, the introduction of dreissenid mussels to upstream locations such as the Luther, Belwood, Guelph, or Conestogo reservoirs would likely lead to the colonization of downstream areas of the Grand River, which are above the Dunnville Dam (Bouvier and Morris 2011). The Sydenham River (both species) and Muskrat Creek (Fawnsfoot only) are the only watersheds that do not contain dreissenid mussels; although, they have been detected at the mouth of the Sydenham River.

Round Goby (*Neogobius melanostomus*), is another invasive species that is spreading throughout the lower Great Lakes and tributaries including the lower Grand, Sydenham and Thames rivers, which may negatively affect both Fawnsfoot and Threehorn Wartyback. Round Goby has been shown to prey on dreissenid mussels (Ghedotti et al. 1995; Ray and Corkum 1997) and has been observed to consume juvenile unionids (M. Poesch, University of Alberta, pers. comm. 2015). It is likely that gape size limitations may be restricting predation on larger mussel species (Ray and Corkum 1997); however, unionids at the juvenile life stage may be vulnerable to consumption due to their smaller size.

Round Goby may also be inhibiting unionid recruitment by acting as a sink for glochidia, meaning that glochidia are unable to survive to the juvenile life stage despite their ability to infest the gills of Round Goby. For example, Tremblay (2012) tested the infestation and metamorphosis rates of four mussel species at risk within the gills of Round Goby and compared them to rates from confirmed host fishes in a laboratory setting. The author concluded that Round Goby serves as a sink for glochidia, not a host, and may be negatively affecting freshwater mussels by disrupting their reproductive cycle (Tremblay 2012).

The host fish(es) for Fawnsfoot and Threehorn Wartyback may potentially be impacted by Round Goby, therefore, it may indirectly affect these populations. Poos et al. (2010) indicate that the potential hosts of Fawnsfoot, Freshwater Drum and Sauger, are not likely to be impacted by Round Goby with regard to competitive interactions; however, a number of studies have documented that Round Goby prey upon the eggs of a wide array of fish species, such as Lake Trout (*Salvelinus namaycush*) (Chotkowski and Marsden 1999), salmonids in general (Fitzsimons et al. 2006), Smallmouth Bass (*Micropterus dolomieu*) (Steinhart et al. 2004), Lake Sturgeon (*Acipenser fulvescens*) (Nichols et al. 2003), and Walleye (*Sander vitreus*) (Roseman et al. 2006), the last of which is a close relative of the Sauger. Taken together, these findings suggest that the potential hosts of Fawnsfoot may be vulnerable to Round Goby predation at earlier life stages.

Another non-indigenous species of great concern to the future of Fawnsfoot and Threehorn Wartyback is the Common Carp (*Cyprinus carpio*), which is thought to be capable of consuming unionids. The feeding behaviours of Common Carp may result in potential harmful habitat alteration, increased turbidity and nutrient levels. At present, Common Carp can be found throughout the distribution of Fawnsfoot and Threehorn Wartyback within the Grand, Sydenham, and Thames rivers (Bouvier et al. 2014). Additional introductions of invasive species into these waters are most likely to occur through the movement of boats from infested areas, use of live baitfish, or natural dispersal of species introduced into the Great Lakes basin.

Turbidity and sediment loading: Increased agricultural land use, which can include clearing of riparian vegetation and unrestricted access to the river by livestock, is often associated with increased sediment loads (WQB 1989a). In addition, increased tile drainage, also related to intensive agricultural practices, often results in large inputs of sediments to the watercourse (COSEWIC 2008). Strayer and Fetterman (1999) noted that increased siltation and suspended solids can affect freshwater mussels by:

- clogging siphons
- inhibiting the intake of oxygen through the clogging gill structures
- reducing the likelihood of the host fish locating the mussel, lure, or conglutinate, as a result of reduced visibility
- reducing flow rates and dissolved oxygen concentrations within the interstitial spaces of the sediment (Österling et al. 2010)

Over 85% and 88% of the Sydenham (DFO 2018a) and lower Thames (Taylor et al. 2004) river watersheds, respectively, are comprised of agricultural land. Suspended solids have been reported as high as 900 mg/L in the Sydenham River (DFO 2018a), a level capable of negatively impacting freshwater mussel populations (Bouvier and Morris 2011), and the lower Thames River is considered to be highly turbid (COSEWIC 2006). Similarly, agricultural and pasture land use, which accounts for 47.8% and 8.3% of the surrounding watershed, respectively, has affected water quality in the Grand River, and the presence of the Dunnville Dam has hindered sediment transport resulting in increased turbidity and sediment loads (Lui et al. 2016; MacDougall and Ryan 2012). It is believed that the greatest impact of this increase will be felt most by species inhabiting the lower portions of the river, such as Fawnsfoot and Threehorn Wartyback (COSEWIC 2006; Bouvier and Morris 2011). The St. Clair River delta is considered to be less affected by this threat, as it is afforded some level of protection (for example, access restrictions) by the Walpole Island First Nation Territory (Bouvier and Morris 2011).

Contaminants and toxic substances: Unionids may be more sensitive to water and sediment contamination (for example, Keller and Zam 1990; Wang et al. 2013) than coexisting fauna. For example, there is evidence suggesting that mussels are sensitive to polychlorinated biphenyls (PCBs), dichlorodiphenyltrichloroethane (DDT), Malathion, and Rotenone; these chemicals can inhibit respiration and accumulate in the tissue of freshwater mussels (U.S. Fish and Wildlife Service [USFWS] 1994). In addition, recent studies have demonstrated that freshwater mussels, at early life stages (glochidia and juveniles), experience greater sensitivity to some contaminants such as ammonia (Augspurger et al. 2003 and 2007), copper (Wang et al. 2007; Gillis et al. 2008 and 2010), and chloride (Gillis 2011; Pandolfo et al. 2012), which are common throughout the watersheds where Fawnsfoot and Threehorn Wartyback occur. In particular, ammonia concentrations have been found to exceed federal guidelines in all sub-basins of the Thames River (Taylor et al. 2004), while the upper range of copper concentrations exceeded provincial water quality objectives in the Grand, Sydenham and Thames rivers (MOECC 1994; Gillis et al. 2010).

These watersheds (Grand, Sydenham, and Thames) in southern Ontario are largely surrounded by agricultural land where activities such as the clearing of riparian zones, the use of fertilizers and pesticides, and the presence of ammonia from tile drainage, wastewater drains, as well as manure storage and spreading, have all contributed to poor water quality. However, in addition to agricultural land use practices, urbanization can also impair water quality. For instance, wild mussels living downstream of a large urban area in the central Grand River exhibited lower condition factor, mean age and elevated levels of stress biomarkers than those living upstream of the cities (Gillis 2012; Gillis et al. 2014). Furthermore, Gillis (2012) observed that concentrations of copper, lead, zinc, aluminum, chromium, and nickel increased significantly in the gills of freshwater mussels continuing downstream in the Grand River, suggesting there is an accumulation of contaminants in downstream areas where Fawnsfoot and Threehorn Wartyback are most likely to occur.

Acidity (Huebner and Pynnonen 1992), and salinity (Liquori and Insler 1985; as cited in USFWS 1994) have also been documented to adversely affect freshwater mussels. The widespread use of road salt (sodium chloride) in the winter months has also been documented to impact freshwater mussel fitness. For example, the glochidia of another unionid species, the Wavyrayed Lampmussel (*Lampsilis fasciola*), have been shown to be highly sensitive to sodium chloride (Gillis 2011). Considering that the ranges of Fawnsfoot and Threehorn Wartyback lie within one of Canada's most road-dense and heavily salted regions, chloride from road salt may pose a significant threat to these species. The current federal water quality guidelines for the protection of aquatic life have been set at 120 mg/L for chronic exposure to chloride, a level that may not sufficiently protect glochidia of some at-risk mussel species (CCME 2011). Further research conducted by Gillis (2011) documented that the upper range of chloride concentrations exceeded the aforementioned water quality guideline within the Grand, Sydenham and Thames rivers. Overall, Todd and Kaltenecker (2012) have conducted studies that suggest long-term road salt use is contributing to increases in baseline chloride concentrations in habitats within the Grand, Sydenham, and Thames rivers where Fawnsfoot and Threehorn Wartyback occur, which may negatively affect the recruitment of these species for years to come.

Various metals and pesticides have been recorded from sediment obtained from the mouths of tributaries to Lake Erie and Lake St. Clair (including areas within the ranges of Fawnsfoot and Threehorn Wartyback) and in concentrations that exceed both federal and provincial standards (Dove et al. 2002, 2003; Bejankiwar 2009). Other concerns include possible endocrine and reproductive effects (for example, Leonard et al. 2014) on freshwater mussels from contaminants contained in municipal effluent. Gagné et al. (2011) determined that the proportion of female unionids (Eastern Elliptio [*Elliptio complanata*]) was significantly higher than males, and that males showed a female-specific protein downstream of a municipal effluent outfall, suggesting that contaminants and toxic substances disrupt gonad physiology and reproduction of this species. Gillis (2012) recorded a negative impact on mussel health (Flutedshell, [*Lasmigona costata*]) and longevity in relation to exposure from urban runoff and municipal wastewater effluents in the Grand River, while Gillis et al. (2014) detected signs of physiological stress as well as possible estrogenic effects in male mussels from short-term exposure (four weeks) to municipal wastewater effluent. The latter study detected elevated concentrations of coliform bacteria, a variety of pharmaceutical and personal care products and natural estrogens at locations downstream of the wastewater plant. Overall, the effects of these contaminants on Fawnsfoot and Threehorn Wartyback are poorly characterized; therefore, further research is needed to elucidate the potential impacts of this threat.

Nutrient loading: Nutrient loading can result from sources including: manure seepage; agricultural runoff; municipal wastewater and sewage discharge; and, riparian clearing and the use of tile drainage practices, which can allow nutrients to infiltrate watersheds more easily. Nutrient loading impacts water quality through eutrophication and leads to increased algal growth and a subsequent reduction of oxygen in the water column (Augsberger et al. 2003). Increased nutrient loads from non-point sources, particularly agricultural sources, were identified by Strayer and Fetterman (1999) as a primary threat to freshwater mussels.

Fawnsfoot and Threehorn Wartyback populations within the Grand, Sydenham, and Thames rivers in southern Ontario are largely surrounded by agricultural land and consequently have been impacted by many of the aforementioned activities (Mackie 1996). Considering that freshwater mussels are highly sensitive to alterations in nutrient levels, these stressors may negatively impact remaining populations. Nutrient loading has been documented within all three of the watersheds where these species are present. For example, the Thames River exhibits some of the highest phosphorus and nitrogen loadings found in the Great Lakes watershed

(WQB 1989b) and phosphorus levels exceeding the provincial water quality objectives are often found in the Sydenham River (DFO 2018a). The Grand River has also received major inputs of nutrients leading to the reclassification of the lower portions as eutrophic-hypereutrophic (MacDougall and Ryan 2012). Within this river, the nitrogen and phosphorous levels are greatest in the spring when increased runoff hastens their transfer from the surrounding agricultural lands into the watershed. These increased inputs of nitrogen and phosphorous have led to a greater prevalence of anoxic areas within the benthic zone during certain times of the year (MacDougall and Ryan 2012), which could potentially kill freshwater mussel species such as Fawnsfoot and Threehorn Wartyback.

Altered flow regimes: Damming of rivers has been shown to detrimentally affect mussels in many ways. Dams alter flow patterns, change the natural thermal profiles of the watercourse, and act as physical barriers to host movement, making large areas of potential habitat completely unavailable to some mussel species (Vaughn and Taylor 1999). Impoundments can result in siltation, stagnation, loss of shallow water habitat, reduced dissolved oxygen, and high concentrations of pollutants and nutrients (Bogan 1993; Vaughn and Taylor 1999; Watters 2000), all of which have been documented in the lower Grand River as a consequence of the Dunnville Dam. Dams can also result in sediment retention upstream and scouring downstream (Bouvier and Morris 2011). There is potential for poor management of water control structures, which may cause de-watering of areas, creating unsuitable habitat for freshwater mussels (Bouvier and Morris 2011) and making them more vulnerable to predation. For more information refer to the RPAs for Fawnsfoot (DFO 2011) and Threehorn Wartyback (DFO 2014).

Habitat removal and alteration: Many activities can result in the physical loss of freshwater mussel habitat. Anthropogenic alterations to the environment, such as channelization, dredging and snagging activities, infilling, and the construction of marinas, docks and impoundments, can have a direct effect on the health of freshwater mussel populations (Watters 2000; Bouvier and Morris 2011). Ultimately, activities that may lead to the increased hardening of substrate in the benthic zone, alterations of the substrate composition, decreased oxygen levels and reduced availability of food resources will have direct impacts on Fawnsfoot and Threehorn Wartyback at multiple life stages.

Availability of host fishes: Any factors that directly or indirectly affect the abundance and distribution of the host fishes will impact the distribution of Fawnsfoot and Threehorn Wartyback. Unionids cannot complete their life cycle without access to the appropriate glochidial host. If host fish populations disappear or decline in abundance to levels below that which can sustain a mussel population, recruitment will no longer occur and the mussel species may become functionally extinct (functionally extinct in this case is defined as a population that is no longer viable, as a crucial part of their life cycle [in this case the host fish] has been removed) (Bogan 1993).

Freshwater Drum, a host fish for Fawnsfoot, may be particularly affected by the presence of dams and barriers considering that its upstream range appears to be limited by the presence of the first dam. This in turn leads to limitations in Fawnsfoot distribution since it is dispersed via the host fish during the glochidial stage. For example, Tiemann et al. (2007) reported that the distribution of Fawnsfoot in the Fox River system of Illinois appears to have been limited by a low-head dam restricting the upstream movement of Freshwater Drum. Within the lower Grand River, Freshwater Drum is currently most abundant downstream of Cayuga, although it is present in lower densities upstream to the Caledonia Dam (MacDougall and Ryan 2012); however, no detections have been made upstream of the dam. Although, Common Shiner and

Longnose Dace, the putative hosts for Threehorn Wartyback, are relatively less dispersive than Freshwater Drum, the presence of dams and barriers also inhibits their upstream dispersal.

Recreational activities: Recreational activities that may impact mussel beds include (Bouvier and Morris 2011):

- driving all-terrain vehicles (ATVs) or other motorized terrestrial vehicles through river beds, this has been identified as a threat in the Sydenham and Thames river
- propellers on recreational boats and personal motorized watercraft; propeller channels have been noted through the mussel beds in the St. Clair River delta
- anglers moving through mussel beds
- paddling action disturbance (kayaks, etc.) of the mussel bed

Recovery

6. Population and distribution objectives

Population and distribution objectives establish, to the extent possible, the number of individuals and/or populations, and their geographic distribution, that is necessary for the recovery of the species. The population and distribution objectives for Fawnsfoot and Threehorn Wartyback are to return self-sustaining populations in the following waterbodies:

1. Grand River (lower)
2. St. Clair River delta (Fawnsfoot only)
3. East Sydenham River
4. North Sydenham River (Bear Creek)
5. Thames River (lower)

The populations at these locations could be considered recovered when they demonstrate active signs of reproduction and recruitment throughout their distribution and are stable or increasing with low risk from known threats. The Great Lakes and connecting channels are specifically excluded from the population and distribution objectives as these areas have been devastated by dreissenid mussels and no longer provide suitable habitat for freshwater mussels. It is currently unknown how long it will take to achieve these objectives but given the threats facing the species and the condition of currently occupied habitat, it is expected to take several decades, if not longer. It should be noted that the setting of population and distribution objectives is a science-based exercise and socio-economic factors were not considered.

Rationale: Very little is known about Fawnsfoot and Threehorn Wartyback populations in Canada and research and monitoring is required before the population and distribution objectives can be refined. For example, population demographics (extent, abundance, trajectories, and targets) are currently unknown.

7. Broad strategies and general approaches to meet objectives

7.1 Actions already completed or currently underway

A monitoring program has been established for the Grand, Sydenham, and Thames rivers. The purpose of these programs was to establish a monitoring network for mussel species at risk throughout the river systems and collect baseline data on their distributions, population demographics and habitat requirements. There are also provisions for the assessment of host fish populations, as well as mussel and host fish habitat monitoring. These programs allow for the tracking of changes in the physical, chemical and biological characteristics of these systems as recovery actions are implemented.

Single- and multi-species recovery strategies have been completed for several freshwater mussel species, the distributions of which partly overlap with Fawnsfoot and Threehorn Wartyback. Recovery teams for these species are currently engaged in the implementation of recovery actions within these watersheds that will benefit Fawnsfoot and Threehorn Wartyback, including:

- Recovery Strategy for the Northern Riffleshell, Snuffbox, Round Pigtoe, Salamander Mussel and Rayed Bean in Canada (DFO 2019)
- Recovery Strategy for the Round Hickorynut (*Obovaria subrotunda*) and the Kidneyshell (*Ptychobranchus fasciolaris*) in Canada (DFO 2013)
- Recovery Strategy and Action Plan for the Mapleleaf (*Quadrula quadrula*) in Canada (Great Lakes-Western St. Lawrence population) (DFO 2018b)

Ecosystem-based recovery initiatives that overlap with Fawnsfoot and Threehorn Wartyback include:

- Sydenham River Action Plan (DFO 2018a), which prescribes the implementation of a number of activities that would benefit Fawnsfoot and Threehorn Wartyback including:
 - investigation into the feasibility of mussel translocations
 - monitoring of mussel and host populations
 - investigation into the impacts of invasive Round Goby
 - continued promotion of mussel identification skills
 - maintenance of flow levels adequate for mussel species
 - establishment of riparian buffer zones in areas of high erosion
 - continue to work with landowners to mitigate the effects of tile drainage and manure waste and advise farm planning and herd management activities
 - public awareness regarding invasive species
- Thames River Ecosystem Recovery Strategy (TRRT 2005). The goal of this strategy involved the development of a “recovery plan that improves the status of all aquatic species at risk in the Thames River through an ecosystem approach that sustains and enhances all native aquatic communities”. Following the lead of the Sydenham River Recovery Team, monitoring stations have also been established in the Thames River.

- Grand River Fish Species at Risk Recovery Strategy (Portt et al. 2007). While this recovery strategy deals specifically with fish species, many of the same threats apply to Fawnsfoot and Threehorn Wartyback, as well as their hosts, such as the impacts of sediment and nutrient loadings and invasive species.
- Walpole Island Ecosystem Recovery Strategy (Bowles 2005)

Other actions that are underway include:

- surveys for unionids, including Fawnsfoot and Threehorn Wartyback, are being conducted as components of research via students and employees from various universities and government agencies, respectively (for example, University of Guelph)
- host fish identification facilities are set up, although the confirmation of the host for Fawnsfoot and Threehorn Wartyback has not occurred

7.2 Measures to be taken to implement the recovery strategy and action plan

Successful recovery of these species is dependent on the actions of many different jurisdictions. It requires the commitment and cooperation of the constituencies that will be involved in implementing the directions and measures set out in this recovery strategy and action plan.

This recovery strategy and action plan provides a description of the measures that provide the best chance of achieving the population and distribution objectives for both Fawnsfoot and Threehorn Wartyback, including measures to be taken to address threats to the species and monitor their recovery, to guide not only activities to be undertaken by DFO, but those for which other jurisdictions, organizations and individuals have a role to play. As new information becomes available, these measures and their priority may change. DFO strongly encourage(s) all Canadians to participate in the conservation of these species by undertaking measures outlined in this recovery strategy and action plan.

Table 7 identifies the measures to be undertaken by DFO to support the recovery of Fawnsfoot and Threehorn Wartyback. Table 8 identifies the measures to be undertaken collaboratively between DFO and its partners, other agencies, organizations or individuals. Implementation of these measures will be dependent on a collaborative approach, in which DFO is a partner in recovery efforts, but cannot implement the measures alone. Table 9 identifies the measures that represent opportunities for other jurisdictions, organizations or individuals. If your organization is interested in participating in one of these measures, please contact the [Ontario and Prairie office](#).

Federal funding programs for species at risk that may provide opportunities to obtain funding to carry out some of the outlined activities include: the [Habitat Stewardship Program for Species at Risk \(HSP\)](#) and the [Aboriginal Fund for Species at Risk Program](#).

While DFO has already commenced efforts to implement the plan, the measures included in this recovery strategy and action plan, but not yet implemented by DFO, will be subject to the availability of funding and other required resources. As indicated in the tables below, partnerships with specific organizations will provide expertise and capacity to carry out some of the listed recovery measures. However, the identification of partners is intended to be advice to

other jurisdictions and organizations and carrying out these actions will be subject to each group's priorities and budgetary constraints.

Four broad strategies were identified to meet the population and distribution objectives: 1) inventory and monitoring; 2) research; 3) communication and outreach; and, 4) stewardship. Approaches are identified for each of these strategies. These approaches are further divided into numbered recovery measures with a priority ranking (high, medium, low); identification of the threat(s) addressed (tables 7 to 9); and, associated timeline (tables 7 and 8). A more detailed narrative is included after the tables (section 7.3). Implementation of the following approaches will be accomplished in coordination with relevant ecosystem-based recovery teams and other pertinent organizations.

Table 7. Measures to be undertaken by Fisheries and Oceans Canada (DFO)

#	Recovery measure	Broad strategy	Approach	Priority ³	Threats or concern addressed	Timeline
1	Conduct/establish routine surveys to monitor changes in the distribution and abundance of current populations as well as invasive species such as dreissenid mussels, Round Goby and Common Carp. Establishing new monitoring stations lower in watersheds where stream reaches are deeper is warranted.	Inventory and monitoring	Assessment	High	Invasive species	Ongoing
2	Conduct further surveys within the historical distribution to detect remnant populations. Determine extent and abundance of any newly discovered remnant populations (that is, North Sydenham River, and Welland River for Fawnsfoot; North Sydenham River and Detroit River for Threehorn Wartyback).	Inventory and monitoring	Assessment	High	Knowledge gaps	Ongoing
3	Conduct targeted surveys in non-historical areas for undetected populations in high probability areas with suitable habitat. Determine extent and abundance of any new populations detected.	Inventory and monitoring	Assessment	Medium	Knowledge gaps	2020 to 2024
4	Establish stations to monitor changes to habitat. This monitoring will inform threat level assessments regarding impacts to sub-	Inventory and monitoring	Assessment	High	Knowledge gaps	Ongoing

³ "Priority" reflects the degree to which the measure contributes directly to the recovery of the species or is an essential precursor to a measure that contributes to the recovery of the species:

- "high" priority measures are considered likely to have an immediate and/or direct influence on the recovery of the species
- "medium" priority measures are important but considered to have an indirect or less immediate influence on the recovery of the species
- "low" priority measures are considered important contributions to the knowledge base about the species and mitigation of threats

#	Recovery measure	Broad strategy	Approach	Priority ³	Threats or concern addressed	Timeline
	populations and be integrated into routine population surveys. It will also allow for the evaluation of progress achieved through recovery implementation activities to reduce threats.					
5	Identify thresholds of tolerance to various threats such as habitat modifications (for example, flow), to determine what constitutes destruction of critical habitat for these two species.	Research	Threat evaluation	High	Turbidity and sediment loading, habitat removal and alteration, altered flow regime	2020 to 2024
6	Characterize the demographic traits of Fawnsfoot and Threehorn Wartyback populations to inform the development of population models.	Research	Life history traits	Medium	Knowledge gaps	2020 to 2025
7	Educate municipal planning authorities so that they consider recovery goals and the protection of critical habitat within official plans.	Communication and outreach	Awareness	Medium	All threats	Ongoing
8	Hold mussel identification workshops that incorporate identification, biology, ecology, threats, and conservation of freshwater mussel species in Ontario.	Communication and outreach	Awareness	Medium	All threats	Ongoing

Table 8. Measures to be undertaken collaboratively between Fisheries and Oceans Canada (DFO) and its partners

#	Recovery measure	Broad strategy	Approach	Priority ⁴	Threats or concern addressed	Timeline (short, medium or long term)	Partner(s) ⁵
9	Determine distribution and abundance of the identified host species (once the functional host has been confirmed) through monitoring and the collection of existing data.	Inventory and monitoring	Assessment	Low	Disruption of host fish(es)	Long term	OMNRF, GRCA, UTRCA, LTVCA, SCRCA, SVCA
10	Determine species sensitivities to environmental contaminants at the glochidial, juvenile and adult life stages.	Research	Threat evaluation	High	Contaminants and toxic substances, sediment and nutrient loading	Medium term	ECCC, MOECC
11	Conduct a risk assessment to identify potential contaminant threats.	Research	Threat evaluation	Medium	Contaminants and toxic substances	Medium term	OMNRF, ECCC, MOECC, GRCA, UTRCA, LTVCA, SCRCA, SVCA
12	Identify physical barriers within the range of the Fawnsfoot and Threehorn Wartyback (determine impact of altered flow on survival).	Research	Threat evaluation	Low	Altered flow regimes	Medium term	OMNRF, GRCA, UTRCA, LTVCA, SVCA

⁴ "Priority" reflects the degree to which the measure contributes directly to the recovery of the species or is an essential precursor to a measure that contributes to the recovery of the species:

- "high" priority measures are considered likely to have an immediate and/or direct influence on the recovery of the species
- "medium" priority measures are important but considered to have an indirect or less immediate influence on the recovery of the species
- "low" priority measures are considered important contributions to the knowledge base about the species and mitigation of threats

⁵ ECCC: Environment and Climate Change Canada; GRCA: Grand River Conservation Authority; LTVCA: Lower Thames Valley Conservation Authority; MOECC: Minister of Environment and Climate Change; OMNRF: Ontario Ministry of Natural Resources and Forestry; SCRCA: St. Clair Region Conservation Authority; SVCA: Saugeen Valley Conservation Authority; UTRCA: Upper Thames River Conservation Authority

#	Recovery measure	Broad strategy	Approach	Priority ⁴	Threats or concern addressed	Timeline (short, medium or long term)	Partner(s) ⁵
13	Investigate the potential impacts of invasive species on host fish abundance.	Research	Threat evaluation	Medium	Invasive species	Medium term	Academia
14	The Sydenham River Action Plan (DFO 2018a) recommended that science-based guidelines should be developed on the feasibility of translocations and repatriations to determine if small populations can be augmented or if the species can be reintroduced in historical ranges. This work would be used to refine population and distribution objectives for Fawnsfoot and Threehorn Wartyback. Artificial propagation would also be beneficial as a means of providing specimens of Fawnsfoot and Threehorn Wartyback that could be used to study the impacts of contaminants.	Research	Population augmentation	Low	Knowledge gaps	Long term	University of Guelph, OMNRF
15	Develop genetically sound propagation guidelines for freshwater mussels, if warranted.	Research	Population augmentation	Low	Knowledge gaps	Long term	University of Guelph, OMNRF
16	Encourage the integration of species recovery and protection into existing watershed plans.	Communication and outreach	Threat mitigation/management	Medium	All threats	Medium term	OMNRF, GRCA, UTRCA, LTVCA, SCRCA

Table 9. Measures that represent opportunities for other jurisdictions, organizations or individuals to lead

#	Recovery measure	Broad strategy	Approach	Priority ⁶	Threats or concern addressed	Suggested other jurisdictions or organizations ⁷
17	Monitor sites where threat mitigation and/or habitat restoration activities have occurred to determine the relative value of recovery measures.	Inventory and monitoring	Assessment	Low	All threats	OMNRF, GRCA, UTRCA, LTVCA, SCRCA
18	Involve local residents, partners, First Nations and appropriate agencies and groups in action planning, habitat improvement and implementation of best management practices (BMPs) to reduce threats.	Communication and outreach	Awareness	High	All threats	GRCA, UTRCA, LTVCA, SCRCA, SVCA, Ontario Federation of Anglers and Hunters (OFAH), OMNRF and First Nations communities
19	Encourage public support and participation by developing awareness materials and programs, which in turn will encourage participation in local stewardship programs and implementation activities to improve and protect habitat.	Communication and outreach	Awareness	Medium	All threats	GRCA, UTRCA, LTVCA, SCRCA, SVCA,
20	Increase awareness within the angling community about the role of hosts for both species (contingent on host fish identification).	Communication and outreach	Awareness	Low	Invasive species, disruption of host fish relationship	GRCA, UTRCA, LTVCA, SCRCA, SVCA, OFAH, OMNRF

⁶ "Priority" reflects the degree to which the measure contributes directly to the recovery of the species or is an essential precursor to a measure that contributes to the recovery of the species:

- "high" priority measures are considered likely to have an immediate and/or direct influence on the recovery of the species
- "medium" priority measures are important but considered to have an indirect or less immediate influence on the recovery of the species
- "low" priority measures are considered important contributions to the knowledge base about the species and mitigation of threats

⁷ GRCA: Grand River Conservation Authority; LTVCA: Lower Thames Valley Conservation Authority; OMNRF: Ontario Ministry of Natural Resources and Forestry; SCRCA: St. Clair Region Conservation Authority; SVCA: Saugeen Valley Conservation Authority; UTRCA: Upper Thames River Conservation Authority

#	Recovery measure	Broad strategy	Approach	Priority ⁶	Threats or concern addressed	Suggested other jurisdictions or organizations ⁷
21	Implement local stewardship programs to improve habitat conditions and reduce threats within critical habitat and other occupied habitats. Priorities and mitigation approaches to be informed through threat evaluation research.	Stewardship	Habitat improvement	High	All threats	GRCA, UTRCA, LTVCA, SCRCA, SVCA,

7.3 Narrative to support the recovery planning and implementation tables

Inventory and monitoring

Recovery measures 1 and 2: A network of monitoring stations has been established for freshwater mussels throughout the current range of Fawnsfoot and Threehorn Wartyback within the East Sydenham River (Metcalf-Smith et al. 2007) and additional monitoring stations using similar methods have more recently been established in the Grand and Thames rivers. Increased monitoring activity will allow for the assessment of the progress made towards achieving the population and distribution objectives prescribed for Fawnsfoot and Threehorn Wartyback. These monitoring sites should be established in a manner so as to permit quantitative tracking of changes in mussel abundance and demographics (size, age, sex), or that of their hosts.

Further surveys are required to confirm the current distribution and derive population estimates for both Fawnsfoot and Threehorn Wartyback in Canada. In addition, monitoring stations should be established in deeper stream reaches of the lower Grand, Sydenham, and Thames rivers to potentially detect Fawnsfoot and Threehorn Wartyback in areas where traditional sampling techniques have been inapplicable, as the majority of their populations are found within these lower reaches. In these cases, new sampling methods involving scuba diving, snorkelling as well as the use of brails⁸ may be warranted to supplement the standard quadrat sampling method designed for shallower habitats. Invasive species monitoring should also be incorporated into the aforementioned freshwater mussel monitoring network. Systems containing both Fawnsfoot and Threehorn Wartyback should be evaluated to determine the current extent of invasions as well as the likelihood of further invasions; plans can be developed to address potential risks, impacts, and proposed actions if monitoring detects the arrival or establishment of invasive species. In addition, further monitoring surveys should also be undertaken within historically occupied areas for both species. For example, further sampling for Fawnsfoot in Lake St. Clair, where dreissenids are less abundant than Lake Erie, as well as the Detroit River where Threehorn Wartyback was recently detected, may confirm that these species are persisting, and enhance estimates of abundance within these waters.

Recovery measure 3: Targeted surveys should also be conducted in watersheds or sub-watersheds where the species were not known to historically occur. These potential areas to be sampled would include areas where the habitat is suitable and the putative host fish species for both Fawnsfoot and Threehorn Wartyback are present, as well as novel areas where fresh shells or live specimens have been discovered. Sampling of the North Sydenham River is recommended for both species as the putative hosts of Fawnsfoot (Freshwater Drum) and Threehorn Wartyback (Common Shiner) are present within the watershed. In addition, another unionid species, Deertoe (*Truncilla truncilla*), has also been recently discovered in a section of the North Sydenham River near Wilkesport. Deertoe is known to use Freshwater Drum as a host in the U.S. (Clark 1981) and exhibits similar habitat preferences as Fawnsfoot, suggesting that this might be a good location to search for the latter species.

⁸ Brails are poles or boards used to drag short chains, with wire prongs that have beaded ends referred to as hooks. When the rig is dragged along a river bed mussels close up as a defense mechanism and consequently clamp down on the beaded ends of the prongs leading to their capture.

Recovery measure 4: Sampling stations should also be established or used to monitor changes in habitat conditions over time. Such habitat assessment would allow for a more thorough evaluation of threats as well as the efficacy of threat mitigation activities implemented to help achieve recovery objectives. These monitoring activities can be led by DFO but can also include monitoring activities conducted by external agencies such as conservation authorities. When combined with population monitoring, habitat tracking can help determine the specific habitat requirements of Fawnsfoot and Threehorn Wartyback as well as the minimum and/or maximum thresholds of tolerance for these species to various threats and water quality parameters (for example, turbidity, contaminant levels, interactions with invasive species). This approach will also assist in the identification of specific areas where habitat restoration or threat mitigation activities should be conducted.

Recovery measure 9: Once the Canadian hosts have been identified, which is a scheduled research activity to refine critical habitat (table 11), it is necessary to determine the distribution, abundance, and health of the host species as they may be a critical factor that limits the abundance and distribution of both Fawnsfoot and Threehorn Wartyback. Due to this relationship, the host species is considered an essential component of the habitat required for freshwater mussel reproduction.

Research

Recovery measures 5, 10 to 14: It is important to evaluate and gain an understanding of how various threats can impact Fawnsfoot and Threehorn Wartyback in order to successfully recover these species. Further research into species-specific tolerance thresholds to various threats will characterize the nature and scale of activities that could be permitted before serious harm to Fawnsfoot and Threehorn Wartyback critical habitat has occurred. Many of the threats facing Fawnsfoot and Threehorn Wartyback can be classified as widespread and chronic (tables 4 and 5) and represent general ecosystem threats that affect numerous other aquatic species. Efforts to remediate these threats will benefit many species in addition to Fawnsfoot and Threehorn Wartyback.

More comprehensive threat evaluations for all extant populations will help inform stewardship programs to ensure the most efficient and effective use of limited resources while promoting an 'ecosystem approach'. Although a recent and thorough RPA has been published for Threehorn Wartyback, threats for Fawnsfoot were evaluated in an older multispecies document; therefore, it is possible that the threat analysis therein may be outdated and less comprehensive. It is also important to investigate potential impacts of contaminants on both Fawnsfoot and Threehorn Wartyback at each specific life stage. Some initial research has been completed on selected contaminants for early life stages of other freshwater mussel species, including chloride, ammonia and copper; however, further work is required that is specific to Fawnsfoot and Threehorn Wartyback along with the watersheds they inhabit. Contaminant levels should also continue to be monitored within the three main watersheds where Fawnsfoot and Threehorn Wartyback occur including the Grand, Sydenham and Thames rivers.

Impoundments may or may not threaten Canadian Fawnsfoot and Threehorn Wartyback populations. Damming of rivers has been shown to detrimentally affect mussels in many ways as it alters flow patterns, changes the natural thermal profiles of the watercourse, and acts as a physical barrier to host species (Vaughn and Taylor 1999). Further research should be undertaken to gain knowledge of how existing dams are affecting Fawnsfoot and Threehorn Wartyback populations to guide mitigation options where required. Evaluations regarding the

feasibility of constructing fish-ways or ladders may be warranted within areas where barriers may be limiting host fish dispersal; however, strong consideration for, and investigation into, the potential for invasive species to spread upstream as a result of such fish passage mechanisms is warranted.

Recovery measure 6: Investigate life-history traits of both the Fawnsfoot and Threehorn Wartyback and characterize the demographics (age at maturation, length at age, fecundity, survivorship, recruitment, etc.) of specific populations to provide the necessary inputs for the development of population models.

Recovery measures 14 and 15: Investigate the potential for population augmentation as a tool for Fawnsfoot and Threehorn Wartyback recovery. This might include the rearing of specimens within aquaculture facilities (stocking), or alternatively, the transfer of wild mussel specimens from a stable donor population (supplementation). Donor populations should be carefully selected to ensure that translocated individuals are likely to experience adequate fitness relative to their new environment. Connected stream sections, that were previously occupied, with the potential to sustain translocated or stocked mussels, will also be identified to increase the likelihood of success and ensure gene flow with other subpopulations.

Communication and outreach

Recovery measures 7, 16, and 18: Development activities that occur outside of identified critical habitat (for example, within the riparian zone or upstream) can still damage or destroy habitat features, particularly when they negatively impact the existing magnitude, timing and frequency of flows. It is important to ensure that planning and management agencies recognize the importance of fluvial processes, the flow and substrate composition requirements of Fawnsfoot and Threehorn Wartyback, as well as the ecological needs of their host fish(es) when reviewing development activities. The protection of critical habitat features should also be considered when developing watershed and sub-watershed management plans. This may also include the removal of obsolete dams in situations where such alteration will not lead to further exposure to threats (for example, upstream spread of invasive species or increased sedimentation). Similarly, consideration for the habitat needs of these species should be communicated to all levels of government to generate support and coordination in terms of protections and recovery measures.

Recovery measures 8 and 20: Increasing freshwater mussel knowledge and identification can be assisted through the development of awareness material, such as the Photo Field Guide for Identifying Freshwater Mussels of Ontario (Metcalf-Smith et al. 2005), and a hands-on workshop that is offered by DFO to government, agency, non-government organizations, Indigenous peoples, the angling community, and the general public. Similarly, the importance of the host fish species to the life cycles of these mussels should also be communicated to the angling community. Increased public knowledge and understanding of the importance of the Fawnsfoot and Threehorn Wartyback, and mussels in general, will play a key role in the recovery of these species. DFO has worked with the Toronto Zoo to develop an online application called Clam Counter for android (available at the Google Play Store) and IOS (available at iTunes) versions.

Recovery measure 19: Conduct outreach sessions and provide information packages that inform the general public of BMPs that landowners can employ to reduce threats to critical habitat. Similarly, this outreach should inform the general public about stewardship and recovery

implementation activities that can be conducted to restore Fawnsfoot and Threehorn Wartyback critical habitat, as well as opportunities for them to become involved as volunteers.

Stewardship

Recovery measure 21: Once threats have been evaluated for extant populations, the results will inform local stewardship programs for threat mitigation. As with other mussels, measures to improve habitat for Fawnsfoot and Threehorn Wartyback include stewardship actions involving BMPs for agricultural properties and residential properties within watersheds where critical habitat has been identified. As Fawnsfoot and Threehorn Wartyback populations are found within the lower portions of the Grand, Sydenham, and Thames river watersheds, it is important to focus stewardship activities within these areas. However, threats including nutrient loading and sedimentation can accumulate from upstream reaches; therefore, the application of BMPs within the upper sections of a watershed may also prove beneficial for Fawnsfoot and Threehorn Wartyback populations. Furthermore, projects funded through HSP including riparian stabilization and planting will further help to improve water quality conditions for both of these mussel species.

8. Critical habitat

8.1 Identification of Fawnsfoot and Threehorn Wartyback critical habitat

8.1.1. General description of Fawnsfoot and Threehorn Wartyback critical habitat

Critical habitat is defined in SARA as "...the habitat that is necessary for the survival or recovery of a listed wildlife species and that is identified as the species' critical habitat in the recovery strategy or in an action plan for the species." [subsection 2(1)]

Also, SARA defines habitat for aquatic species as "...spawning grounds and nursery, rearing, food supply, migration and any other areas on which aquatic species depend directly or indirectly in order to carry out their life processes, or areas where aquatic species formerly occurred and have the potential to be reintroduced." [subsection 2(1)]

For Fawnsfoot and Threehorn Wartyback, critical habitat is identified to the extent possible, using the best available information, and provides the functions and features necessary to support the species' life-cycle processes and to achieve the species' population and distribution objectives.

This recovery strategy and action plan identifies critical habitat for Fawnsfoot and Threehorn Wartyback as reaches with moderate to low flows, and mud to sand or fine gravel substrates within the Grand, Sydenham, and Thames rivers.

The critical habitat identified for Fawnsfoot in this recovery strategy and action plan is insufficient to achieve the population and distribution objectives as critical habitat has not been identified within the St. Clair Delta where the species historically and currently occurs. It is unknown if the critical habitat identified for the Threehorn Wartyback in this recovery strategy and action plan is sufficient to achieve the species' population and distribution objectives. The schedule of studies in section 8.2 outlines the research required to acquire more detailed information about the critical habitat identified to achieve the species' population and distribution

objectives as the species may be more widely distributed within the watersheds where it is found. In addition, new information characterizing the functions, features and attributes of critical habitat for both species may also be refined.

8.1.2. Information and methods used to identify critical habitat

Within the rivers currently occupied by Fawnsfoot and Threehorn Wartyback, an ecological classification system was used, along with species detections and applicable sampling effort, to identify critical habitat. The OMNRF's Aquatic Landscape Inventory System (ALIS version 1) (Stanfield and Kuyvenhoven 2005) was used as the base unit for defining reaches within riverine systems. The ALIS system employs a valley classification approach to define river segments with similar habitat and continuity on the basis of hydrography, surficial geology, slope, position, upstream drainage area, climate, land cover and the presence of instream barriers, all of which are believed to have a controlling effect on the biotic and physical processes within the catchment. Therefore, if the species have been found in one part of the ecological classification, it would be reasonable to expect that they would be present in other spatially contiguous areas of the same valley segment. Within all identified river segments (that is, valley segments), the width of the habitat zone is defined as the area from the mid-channel point to bankfull⁹ width on both the left and right banks; this supports long-term channel forming discharges important in maintaining in-stream habitat conditions required by freshwater mussels. Critical habitat for Fawnsfoot and Threehorn Wartyback was therefore identified as the reach of river that includes all contiguous ALIS segments from the uppermost stream segment with the species present to the lowermost stream segment with the species present; segments or reaches were excluded only when supported by robust data indicating the species' absence and/or unsuitable habitat conditions. Current occupancy for these species was defined by recent records of live individuals (and/or fresh shells) from 1996 onward; this is the point in time when systematic surveys of freshwater mussel communities in southern Ontario began. Unoccupied ALIS segments with suitable habitats were also included when limited sampling had occurred (that is, the species were assumed to be present).

8.1.3. Identification of critical habitat

Geographic information

For Fawnsfoot and Threehorn Wartyback, critical habitat is identified in the Grand, East Sydenham, North Sydenham (Bear Creek), and Thames rivers (figures 7 to 14); additional areas of potential critical habitat within the St. Clair River delta region will be considered in collaboration with Walpole Island First Nation. Critical habitat has not been identified within Rondeau Bay since both species are considered to be extirpated from this location as a result of the invasion of dreissenid mussels and the lack of detections within the last 20 years. The locations of the critical habitat's functions, features and attributes have been identified using the 'bounding box' approach. This means that the critical habitat is not comprised of the entire area within the identified boundaries but only those areas within the identified geographical boundaries (table 10) where the biophysical feature and the function it supports occur, as described in table 11.

⁹ From the top of the riverbank on one side of the channel to the top of the riverbank on the other.

Table 10. Coordinates locating the boundaries within which critical habitat is found for Fawnsfoot and Threehorn Wartyback

Location	Point 1	Point 2
Grand River (both species)	42.8569, -79.5769	43.0732, -79.9589
North Sydenham River (Bear creek) (both species)	42.7285, -82.3511	42.8796, -82.1394
East Sydenham River/Bear Creek (both species)	42.5612, -82.4117	42.7669, -81.8473
Thames River (Fawnsfoot)	42.3185, -82.4539	42.9771, -81.3707
Thames River (Threehorn Wartyback)	42.5266, -82.0382	42.9771, -81.3707

North Sydenham River (Bear Creek): The area within which critical habitat is currently identified is the ALIS segments with the species present (figures 7 and 11). For both Fawnsfoot and Threehorn Wartyback, this represents a total river reach of approximately 50 km in length and is described as follows:

- beginning on Bear Creek approximately 200 m downstream of Petrolia Line at the Bridgeview Park Dam in Petrolia and continuing downstream to a point approximately 220 m upstream of Kimball Road where Bear Creek and Black Creek join to form the North Sydenham River

East Sydenham River: The area within which critical habitat is currently identified is a partial ALIS segment with the species present; however, critical habitat is terminated in this segment due to the fact that upstream reaches within this segment have been heavily sampled and no Fawnsfoot or Threehorn Wartyback have been detected (figures 8 and 12). For both the Fawnsfoot and Threehorn Wartyback, this represents a total river reach of approximately 100 km in length and is described as follows:

- beginning downstream of Cameron Road between Sydenham Line and Aughrim Line, and continuing downstream to the Sydenham River mouth at the St. Clair River delta

Thames River: The area within which critical habitat is currently identified is the ALIS segments with the species present (figures 9a, 9b, and 13). For Fawnsfoot, this represents a total river reach of approximately 212 km in length and is described as follows:

- beginning approximately 2.5 km upstream of the Oxford Street West bridge in Kilworth in an area adjacent to Kains Woods Environmental Area and continuing downstream to the Thames River mouth at Lake St. Clair in Lighthouse Cove (north of Tilbury)

For Threehorn Wartyback this represents a total river reach of approximately 116 km and can be described as follows:

- beginning approximately 2.5 km upstream of the Oxford Street West bridge in Kilworth in an area adjacent to Kains Woods Environmental Area and continuing downstream to a point approximately 14 km downstream of Victoria Road, and approximately 0.75 km south of Longwoods Road (Highway 2)

Grand River: The area within which critical habitat is currently identified is the ALIS segments with the species present (figures 10 and 14). For both species, this represents a total river reach of approximately 70 km in length and is described as follows:

- beginning immediately below the Caledonia Dam (Grand River Dam) and continuing downstream to the Grand River mouth at Lake Erie in Port Maitland




Figure 7. Area within which critical habitat is found for Fawnsfoot in the North Sydenham River (Bear Creek)




Figure 8. Area within which critical habitat is found for Fawnsfoot in the East Sydenham River




Figure 9a. Area within which critical habitat is found for Fawnsfoot in the lower Thames River (east)




Figure 9b. Area within which critical habitat is found for Fawnsfoot in the lower Thames River (west)




Figure 10. Area within which critical habitat is found for Fawnsfoot in the Grand River




Figure 11. Area within which critical habitat is found for Threehorn Wartyback in the North Sydenham River (Bear Creek)




Figure 12. Area within which critical habitat is found for Threehorn Wartyback in the East Sydenham River




Figure 13. Area within which critical habitat is found for Threehorn Wartyback in the Thames River




Figure 14. Area within which critical habitat is found for Threehorn Wartyback in the Grand River

Biophysical functions, features and attributes

Table 11 summarizes the best available knowledge of the functions, features and attributes for each life stage of the Fawnsfoot and Threehorn Wartyback (refer to section 4.3 needs of the species for full references). Note that not all attributes in table 11 must be present for a feature to be identified as critical habitat. If the features, as described in table 11, are present and capable of supporting the associated function(s), the feature is considered critical habitat for the species, even though some of the associated attributes might be outside of the range indicated in the table.

Table 11. Essential functions, features and attributes of critical habitat for each life stage of the Fawnsfoot and Threehorn Wartyback (riverine populations)¹⁰

Species	Life stage	Function ¹¹	Feature(s) ¹²	Attribute(s) ^{13, 14}
Fawnsfoot	Spawning and fertilization (time period unknown) Glochidia present in females (long-term brooder: spring/summer, spring)	Reproduction	Lakes and reaches of rivers and streams with moderate to low flows, mud to sand or fine gravel substrates (includes 'bankfull channel' ¹⁵) and suitable water quality parameters	Attributes assumed to be same as for adults (see below): 1. flow present (distribution of sperm) 2. low contaminants levels below the following: a) long-term chloride levels < 120 mg/L (CCME 2011) b) mean concentrations of < 0.3 mg/L total ammonia as Nitrogen at pH 8; for protection of all life stages of freshwater mussels (Augsburger et al. 2003) c) copper levels < 3 µg/L (CCME 2005) should protect sensitive glochidia (Gillis et al. 2008)

¹⁰ Where known or supported by existing data.

¹¹ Function: life-cycle process of the listed species taking place in critical habitat (for example, spawning, nursery, rearing, feeding and migration). The function informs the rationale for its protection. The identification of critical habitat must describe how the functions support a life process necessary for the survival or recovery of species at risk.

¹² Feature: every function is the result of a single or multiple feature(s), which are the structural components of the critical habitat. Features describe how the habitat is critical and they are the essential structural component that provides the requisite function(s) to meet the species' needs. Features may change over time and are usually comprised of more than one part, or attribute. A change or disruption to the feature or any of its attributes may affect the function and its ability to meet the biological needs of the species.

¹³ Attribute: Attributes are measurable properties or characteristics of a feature. Attributes describe how the identified features support the identified functions necessary for the species' life processes. Together, the attributes allow the feature to support the function. In essence, attributes provide the greatest level of information about a feature, the quality of the feature and how the feature is able to support the life-cycle requirements of the species.

¹⁴ Note that not all attributes must be present for a feature to be identified as critical habitat.

¹⁵ From the top of the riverbank on one side of the channel to the top of the riverbank on the other.

Species	Life stage	Function ¹¹	Feature(s) ¹²	Attribute(s) ^{13, 14}
Threehorn Wartyback	Spawning and fertilization (short-term brooder: glochidia forming and being released in May until the end of July)	Same as above	Same as above	Same as above
Both species	Encysted glochidial stage (time period unknown) on host fish(es) until drop off	Development on host	Same as above with host fish(es) present	Attributes assumed to be same as below (as these conditions support both host fish[es] and adults): 3. presence of host fish(es), Fawnsfoot, Freshwater Drum and Sauger; Threehorn Wartyback, Common Shiner and Longnose Dace ¹⁶ 4. dissolved oxygen [DO] levels sufficient to support host(s) (> 4 mg/L [MOECC 1994] for protection of warm water species)
Fawnsfoot	Adult/juvenile	Feeding, cover, nursery	Same as above	5. Usually sand or mud (Clark 1981); has been found in areas with coarser substrates (Howells et al. 1996) 6. Depths from 1 to > 5 m
Threehorn Wartyback	Adult/juvenile	Feeding, cover, nursery	Same as above	7. Most often found in areas where the substrate is composed of boulder, rubble, gravel and sand, or some combination thereof
Both species	Adult/juvenile	Feeding, cover, nursery	Same as above	8. Adequate supply of food (plankton: bacterial, algae, organic detritus, protozoans) 9. Dreissenids absent or in low abundance 10. Maintenance of an “environmental thermal regime” ¹⁷ (gamete production and development) 11. No threat of harm and harassment to individuals

¹⁶ Host fish(es) for both Fawnsfoot and Threehorn Wartyback have yet to be confirmed in a laboratory setting.

¹⁷ Maintenance of an ‘environmental thermal regime’ requires that water temperatures are maintained within the limits of natural variability (daily or seasonal) such that life-cycle processes are completed without impacting the fitness of the organism.

Studies to further refine knowledge on the essential functions, features and attributes for various life stages of the Fawnsfoot and Threehorn Wartyback are described in section 8.2 (schedule of studies to identify critical habitat).

Summary of critical habitat relative to population and distribution objectives

These are areas that, based on current best available information, the Minister of Fisheries and Oceans considers necessary to partially achieve the species’ population and distribution objectives required for the survival/recovery of these two species. Additional critical habitat may be identified in future updates to the recovery strategy and action plan.

8.2 Schedule of studies to identify critical habitat

Further research is required to refine the understanding of the functions, features and attributes of the currently identified critical habitat necessary to support the species’ population and distribution objectives and protect the critical habitat from destruction. The activities listed in table 12 are not exhaustive and it is likely that the process of investigating these actions will lead to the discovery of further knowledge gaps that need to be addressed.

Table 12. Schedule of studies to refine critical habitat for both Fawnsfoot and Threehorn Wartyback

Description of activity	Rationale	Timeline*
Determine habitat requirements of all life stages (glochidia, juvenile, and adult) of the Fawnsfoot and Threehorn Wartyback as well as the host fish species (once the functional hosts have been confirmed).	Refine features and attributes of critical habitat and determine if unique conditions are required for any particular life stage.	2020 to 2024
Determine/confirm functional host fish species.	Will determine hosts for the glochidia (parasitic larvae) to juvenile transformation.	2020 to 2024
Based on collected information, review population and distribution objectives. Determine amount (potential for further critical habitat to be identified), configuration and description of critical habitat required to achieve these objectives if adequate information exists.	Refinement of recovery objectives as well as critical habitat description to meet these objectives.	Ongoing

* Timelines are subject to change in response to demands on resources and/or personnel and as new priorities arise.

8.3 Activities likely to result in the destruction of critical habitat

The following examples of activities likely to result in the destruction¹⁸ of critical habitat (table 13) are based on known human activities that are likely to occur in and around critical habitat and would result in the destruction of critical habitat if unmitigated. The list of activities is neither exhaustive nor exclusive and has been guided by the threats described in section 5 (threats). The absence of a specific human activity from this table does not preclude or restrict DFO’s

¹⁸ Destruction occurs when there is a temporary or permanent loss of a function of critical habitat at a time when it is required by the species.

ability to regulate that activity under SARA. Furthermore, the inclusion of an activity does not result in its automatic prohibition, and does not mean the activity will inevitably result in destruction of critical habitat. Every proposed activity must be assessed on a case-by-case basis and site-specific mitigation will be applied where it is reliable and available. Where information is available, thresholds and limits have been developed for critical habitat attributes to better inform management and regulatory decision making. However, in many cases, knowledge of a species and its critical habitat's thresholds of tolerance to disturbance from human activities is lacking and must be acquired.

Table 13. Examples of human activities likely to result in the destruction of critical habitat for both Fawnsfoot and Threehorn Wartyback. The pathway of effect for each activity is provided as well as the potential links to the biophysical functions, features, and attributes of critical habitat

Threat	Activity ¹⁹	Effect pathway	Life stage affected	Function affected	Feature affected	Attribute affected ²⁰
Turbidity and sediment loading	<p>Work in or around water with improper sediment and erosion control (for example, installation of bridges, pipelines, culverts, overland runoff from ploughed fields, run-off from urban and residential development, use of industrial equipment, cleaning or maintenance of bridges or other structures without proper mitigation).</p> <p>Unfettered livestock access to waterbodies.</p> <p>Removal or cultivation of riparian vegetation.</p>	<p>Improper sediment and erosion control or mitigation can cause increased turbidity and sediment deposition, changing preferred substrates and impairment of feeding and reproductive functions.</p> <p>When livestock have unfettered access to waterbodies, damage to shorelines, banks and watercourse bottoms can cause increased erosion and sedimentation, affecting turbidity and water temperatures.</p> <p>Agricultural lands, particularly those with little riparian vegetation and without tile drainage, allow large inputs of sediments into the watercourse.</p>	All	All	Reaches of rivers and streams with mud to sand or fine gravel substrates, suitable water quality parameters and host fish present	2 (a, b, c), 3, 4, 5, 6, 7, 8, 10
Nutrient loading	Over-application of fertilizer and improper nutrient management (for example, organic debris management, wastewater management, animal waste, septic systems and municipal sewage).	<p>Improper nutrient management can cause nutrient loading of nearby waterbodies. Elevated nutrient levels (phosphorus and nitrogen) can cause increased turbidity causing harmful algal blooms, changing water temperatures, and reduced DO levels.</p> <p>Mussel survival rates are closely related to DO levels. Low DO may</p>	All	All	Reaches of rivers and streams with suitable water quality parameters	2(b), 3, 4, 8, 10

¹⁹ These activities affect multiple life stages, and therefore, multiple functions and potentially multiple features.

²⁰ Please refer to table 11 (essential functions, features and attributes of critical habitat for each life stage of the Fawnsfoot and Threehorn Wartyback) for details regarding attributes associated with specific life stages.

Threat	Activity ¹⁹	Effect pathway	Life stage affected	Function affected	Feature affected	Attribute affected ²⁰
		<p>cause mortality of host fishes, thereby disrupting mussel reproductive cycles.</p> <p>Recent evidence has shown that juvenile mussels are among the most sensitive aquatic organisms to ammonia toxicity.</p>				
Altered flow regimes	<p>Change in timing, duration and frequency of flow. Water-level management (for example, through dam operation) or water extraction activities (for example, for irrigation), that causes dewatering of habitat or excessive flow rates. Similarly, this threat includes the hastened discharge from surrounding watersheds due to large increases in impervious surfaces from urban and residential development.</p>	<p>High flow conditions (and 'flashier' flows) can cause dislodgement and passive transport of mussels from areas of suitable habitat into areas of lesser or marginal habitat.</p> <p>Low flows can result in depressed DO levels, elevated temperatures and stranding. Host fishes may also be impacted, thereby disrupting reproduction.</p> <p>Altered flow patterns can affect habitat availability (for example, by 'dewatering' habitats) in creeks and rivers, sediment deposition (for example, changing preferred substrates), and water temperatures.</p>	All	All	Reaches of rivers and streams with moderate to low flows and mud to sand or fine gravel substrates	1, 3, 4, 5, 6, 7, 8, 9, 10
Contaminants and toxic substances	<p>Over application or misuse of herbicides and pesticides.</p> <p>Release of urban runoff and municipal and industrial pollution into habitat (including municipal wastewater effluents).</p>	<p>Introduction of toxic compounds (for example, high chloride levels, pharmaceutical pollutants. from stormwater runoff) into habitat used by these species can change water chemistry affecting the habitat and host fish availability or use, especially during sensitive life stages.</p>	All	All	Reaches of rivers and streams with suitable water quality parameters	2 (a, b, c), 3, 4, 8

Threat	Activity ¹⁹	Effect pathway	Life stage affected	Function affected	Feature affected	Attribute affected ²⁰
	Introduction of high levels of chloride through activities such as excessive salting of roads in winter.	Chloride levels have shown recent inclines due to an increase in the use of road salt. High chloride levels can cause direct mortality of sensitive glochidia.				
Habitat removal and alteration	<ul style="list-style-type: none"> - Dredging - Grading - Excavation <p>Placement of material or structures in water (for example, pilings, bridge construction, infilling, partial infills), construction of dams and barriers, as well as shoreline hardening.</p>	<p>Changes in bathymetry, shoreline and channel morphology caused by dredging and nearshore grading and excavation can alter preferred substrates, change water depths, change flow patterns potentially affecting turbidity, nutrient levels and water temperatures.</p> <p>Placing material or structures in water reduces habitat availability (for example, the footprint of the infill or structure is lost). Placing of fill can cover preferred substrates for mussels and their host fishes.</p> <p>Dams/barriers can result in direct loss of habitat or fragmentation, which can limit the reproductive capabilities and dispersal of mussels by eliminating or decreasing the number of hosts available as well as host dispersal.</p> <p>Hardening of shorelines can reduce organic inputs into the water and alter water temperatures, potentially affecting the availability of food for these species.</p>	All	All	Reaches of rivers and streams with moderate to low flows and mud to sand or fine gravel substrates, as well as suitable water quality parameters	1, 3, 4, 5, 6, 7, 8, 10
Recreational activities	<p>Baitfish releases.</p> <p>Use of motor vehicles in river.</p>	<p>Spread aquatic invasive species (boats, bait buckets).</p> <p>Disrupt substrate, dislodge mussels.</p>	All	All	Reaches of rivers and streams with moderate to low flows and mud to	3, 4, 8, 11

Threat	Activity ¹⁹	Effect pathway	Life stage affected	Function affected	Feature affected	Attribute affected ²⁰
					sand or fine gravel substrates, as well as suitable water quality parameters	
Disruption of host fishes	Excessive removal of host fishes (through either commercial or recreational harvest) or indirect means (for example, damming activities may prevent fish movement).	Any activities that affect the host species' abundance, movements, or behaviour during the period of encystment or release may disrupt the reproductive cycle of these mussels.	Encysted glochidial stage	Development on hosts	Reaches of rivers and streams with moderate to low flows, mud to sand or fine gravel substrates, suitable water quality parameters, and host fish(es) present	3

In future, threshold values for some stressors may be informed through further research. For some of the above activities, BMPs may be enough to mitigate threats to the species and its habitat; however, in some cases, it is not known if BMPs are adequate to protect critical habitat or if further research is required.

8.4 Proposed measures to protect critical habitat

Under SARA, critical habitat must be legally protected from destruction within 180 days of being identified in a final recovery strategy or action plan and included in the Species at Risk Public Registry. For the critical habitat of both Fawnsfoot and Threehorn Wartyback, it is anticipated that this will be accomplished through a SARA Critical Habitat Order made under subsections 58(4) and (5), which will invoke the prohibition in subsection 58(1) against the destruction of the identified critical habitat.

9. Evaluation of the socio-economic costs and of benefits of the action plan

SARA requires that the action plan component of the recovery document²¹ (action plan) include an evaluation of the socio-economic costs of the action plan and the benefits to be derived from its implementation (SARA 49(1)(e), 2003). This evaluation addresses only the incremental socio-economic costs of implementing this action plan from a national perspective as well as the social and environmental benefits that would occur if the action plan were implemented in its entirety, recognizing that not all aspects of its implementation are under the jurisdiction of the federal government. It does not address cumulative costs of species recovery in general nor does it attempt a cost-benefit analysis. Its intent is to inform the public and to guide decision making on implementation of the action plan by partners.

The protection and recovery of species at risk can result in both benefits and costs. The Act recognizes that “wildlife, in all its forms, has value in and of itself and is valued by Canadians for aesthetic, cultural, spiritual, recreational, educational, historical, economic, medical, ecological and scientific reasons” (SARA 2003). Self-sustaining and healthy ecosystems with their various elements in place, including species at risk, contribute positively to the livelihoods and the quality of life of all Canadians. A review of the literature confirms that Canadians value the preservation and conservation of species in and of themselves. Actions taken to preserve a species, such as habitat protection and restoration, are also valued. In addition, the more an action contributes to the recovery of a species, the higher the value the public places on such actions (Loomis and White 1996; DFO 2008). Furthermore, the conservation of species at risk is an important component of the Government of Canada’s commitment to conserving biological diversity under the International Convention on Biological Diversity. The Government of Canada has also made a commitment to protect and recover species at risk through the Accord for the Protection of Species at Risk. The specific costs and benefits associated with this action plan are described below.

It is important to note that the socio-economic evaluation only applies to the detailed recovery measures. The setting of population and distribution objectives and the identification of critical habitat are science-based exercises and socio-economic factors were not considered in their development.

This evaluation does not address the socio-economic impacts of protecting critical habitat for Fawnsfoot and Threehorn Wartyback. Under SARA, DFO must ensure that critical habitat identified in a recovery strategy or action plan is legally protected within 180 days of the final posting of the recovery strategy or action plan. Where a Critical Habitat Order will be used for critical habitat protection, the development of the Order will follow a regulatory process in

²¹ That is, tables 7 to 9 and section 9

compliance with the Cabinet Directive on Regulatory Management, including an analysis of any potential incremental impacts of the SARA Order that will be included in the Regulatory Impact Analysis Statement. As a consequence, no additional analysis of the critical habitat protection has been undertaken for the assessment of costs and benefits of the action plan.

Policy baseline

The policy baseline consists of the protection under SARA for Fawnsfoot and Threehorn Wartyback (both species were listed under SARA in 2019), along with continued protection under Ontario's *Endangered Species Act, 2007*. Other legislation that may provide direct or indirect habitat protection for these species includes the federal *Fisheries Act* and existing provincial legislation.

The policy baseline also includes any recovery measures that were implemented prior to and after Fawnsfoot and Threehorn Wartyback were listed under SARA. These recovery measures include recovery strategies and action plans for other freshwater species as well as multispecies ecosystem-based recovery programs discussed in section 7.1 of this report.

Socio-economic costs

The majority of the recovery activities identified in this recovery strategy and action plan are short term (2020 to 2024), medium term, or ongoing. Most of these activities focus on research, monitoring, engagement, education, stewardship and outreach, and management to reduce threats and to inform and promote species recovery. Some of the actions are one-time projects likely funded from existing federal government resources. Implementation of local stewardship actions would be supported by programs such as HSP. In addition, most programs require a level of direct or in-kind support costs from applicants as matching funds. The costs (direct and in-kind) associated with these short-term actions are estimated to be low and spread over the next five years.

Costs would be incurred by the federal government and its partners to implement the activities listed in the recovery strategy and action plan. In-kind costs, such as volunteer time, providing expertise and equipment, would be incurred as a result of implementing activities listed in the recovery strategy and action plan. Costs (including in-kind support) could be incurred by the province of Ontario and conservation authorities. Land owners along the waterways where Fawnsfoot and Threehorn Wartyback are present may incur some costs for BMPs. However, as many of the activities and actions are implemented on a collaborative and voluntary nature, land owners are likely to only incur costs on a voluntary basis.

Long-term recovery activities will be developed through a cooperative approach following discussions between other agencies, levels of government, stewardship groups and stakeholders allowing for consideration of costs and benefits during the process.

Socio-economic benefits

Some of the benefits of recovery actions required to return self-sustaining populations of Fawnsfoot and Threehorn Wartyback outlined in this recovery strategy and action plan are difficult to quantify but would generally be positive. Freshwater mussels play an integral role in the functioning of aquatic ecosystems. They are sensitive indicators (for example, water and habitat quality) of the health of freshwater ecosystems. These ecosystem benefits would be

maintained as a result of implementing the recovery actions proposed in the recovery strategy and action plan.

Some of the unquantifiable non-market benefits mentioned in the second paragraph of this evaluation would be enjoyed by the Canadian public as a result of implementing the recovery actions contained in the recovery strategy and action plan. Recovery actions that improve riverine habitat will help lead to healthier watersheds with benefits such as improved water quality.

The benefits of implementing the recovery actions contained in the recovery strategy and action plan are anticipated to be low.

Distributional impacts

Governments (federal and provincial) and conservation authorities will incur the majority of costs of implementing the recovery strategy and action plan.

The Canadian public will benefit from the implementation of the recovery strategy and action plan through expected non-market and ecosystem benefits associated with recovery and protection of the species and their habitats. Recovery actions that improve riverine habitat will help lead to healthier watersheds with benefits such as improved water quality.

10. Measuring progress

The performance indicators presented below provide a way to define and measure progress toward achieving the population and distribution objectives. A successful recovery program will achieve the overall aim of recovering populations to a state where they are stable or increasing with low risk from known threats. Progress towards meeting these objectives will be reported on in the report on the progress of recovery strategy implementation.

Reporting on the implementation of the action plan components, under section 55 of SARA, will be completed by assessing the degree of progress towards achieving the broad strategies/approaches outlined in this document. Reporting on the ecological and socio-economic impacts of the action plan, under section 55 of SARA, will be completed by assessing:

- a) the results of monitoring the recovery of the species
- b) the species' long-term viability; and
- c) the implementation of the action plan.

Performance indicators:

1. The continued presence of Fawnsfoot and Threehorn Wartyback within their known distribution by 2024
2. Confirmed reproduction at known sites by 2024

11. References

- Augspurger, T., F.J. Dwyer, C.G. Ingersoll, C.M. Kane. 2007. Advances and opportunities in assessing contaminant sensitivity of freshwater mussel (Unionidae) early life stages. *Environmental Toxicology and Chemistry* 26: 2025–2028.
- Augspurger, T., A.E. Keller, M.C. Black, W.D. Cope, and F.J. Dwyer. 2003. Water quality guidance for protection of freshwater mussels (Unionidae) from ammonia exposure. *Environmental Toxicology and Chemistry* 22: 2569–2575.
- Baker, S.M. and D.J. Hornbach. 1997. Acute physiological effects of Zebra Mussel (*Dreissena polymorpha*) infestation on two unionid mussels, *Actinonaias ligamentina* and *Amblema plicata*. *Canadian Journal of Fisheries and Aquatic Sciences* 54: 512–519.
- Barnhart, M.C. and M.S. Baird. 2000. Fish hosts and culture of mussel species of special concern: Annual report for 1999. U.S. Fish and Wildlife Service and Natural History Section, Missouri. 39 p.
- Bejankiwar, R. 2009. Essex Region Conservation Authority: Water quality status report. Accessed 12 December 2012). 43 p.
- Bogan, A.E. 1993. Freshwater bivalve extinctions (Mollusca: Unionoida): a search for causes. *American Zoologist* 33: 599–609.
- Bouvier, L.D. and T.J. Morris. 2011. Information in support of a recovery potential assessment of Eastern Pondmussel (*Ligumia nasuta*), Fawnsfoot (*Truncilla donaciformis*), Mapleleaf (*Quadrula quadrula*), and Rainbow (Villose iris) in Canada. DFO Canadian Science Advisory Secretariat Research Document 2010/120. vi + 51 p.
- Bouvier L.D., J.A.M. Young, and T.J. Morris. 2014. Information in support of a recovery potential assessment of Threehorn Wartyback (*Obliquaria reflexa*) in Canada. DFO Canadian Science Advisory Secretariat Research Document 2014/023. v + 38 p.
- Bowles, J.M. 2005. Walpole Island ecosystem recovery strategy (Draft 8). Prepared for the Walpole Island Heritage Centre, Environment Canada and the Walpole Island Recovery Team. 45 p.
- CCME (Canadian Council of Ministers of the Environment). 2005. Canadian water quality guidelines. Canadian Council of Ministers of the Environment, Environment Canada, Ottawa, ON.
- CCME (Canadian Council of Ministers of the Environment). 2011. Canadian water quality guidelines (chloride). Canadian Council of Ministers of the Environment, Environment Canada, Ottawa, ON.
- Chotkowski, M.A. and J.E. Marsden. 1999. Round Goby and Mottled Sculpin predation on Lake Trout eggs and fry: field predictions from laboratory experiments. *Journal of Great Lakes Research* 25: 26–35.

Clarke, A.H. 1981. The Freshwater Molluscs of Canada. National Museums of Canada, Ottawa. 446 p.

COSEWIC (Committee on the Status of Endangered Wildlife in Canada). 2006. COSEWIC assessment and status report on the Mapleleaf mussel, *Quadrula quadrula* (Saskatchewan - Nelson population and Great Lakes - Western St. Lawrence population) in Canada. vii + 58 p.

COSEWIC (Committee on the Status of Endangered Wildlife in Canada). 2008. COSEWIC assessment and status report on the Fawnsfoot, *Truncilla donaciformis*, in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa. vii + 39 p. ([SAR Public Registry Fawnsfoot](#)) (Accessed: October, 2018).

COSEWIC (Committee on the Status of Endangered Wildlife in Canada). 2013. COSEWIC assessment and status report on the Threehorn Wartyback *Obliquaria reflexa* in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa. ix + 58 p. ([SAR Public Registry Threehorn Wartyback](#)) (Accessed: October, 2018).

DFO (Fisheries and Oceans Canada). 2008. Estimation of the economic benefits of marine mammal recovery in the St. Lawrence estuary. Policy and Economics Regional Branch, Quebec 2008.

DFO (Fisheries and Oceans Canada). 2011. Recovery potential assessment of Eastern Pondmussel (*Ligumia nasuta*), Fawnsfoot (*Truncilla donaciformis*), Mapleleaf (*Quadrula quadrula*), and Rainbow (Villose iris) in Canada. DFO Canadian Science Advisory Secretariat Science Advisory Report 2010/073.

DFO (Fisheries and Oceans Canada). 2013. Recovery strategy for the Round Hickorynut (*Obovaria subrotunda*) and Kidneyshell (*Ptychobranthus fasciolaris*) in Canada. In *Species at Risk Act Recovery Strategy Series*, Fisheries and Oceans Canada, Ottawa. vi + 70 p.

DFO (Fisheries and Oceans Canada). 2014. Recovery Potential Assessment of Threehorn Wartyback (*Obliquaria reflexa*). DFO Canadian Science Advisory Secretariat Science Advisory Report 2014/014.

DFO (Fisheries and Oceans Canada). 2018a. Action plan for the Sydenham River in Canada: an ecosystem approach. In *Species at Risk Act Action Plan Series*. Fisheries and Oceans Canada, Ottawa. iv + 35 p.

DFO (Fisheries and Oceans Canada). 2018b. Recovery strategy and action plan for the Mapleleaf (*Quadrula quadrula*) in Canada (Great Lakes-Western St. Lawrence population) [Proposed]. *Species at Risk Act Recovery Strategy Series*. Fisheries and Oceans Canada, Ottawa. vi + 57 p.

DFO (Fisheries and Oceans Canada). 2019. Recovery strategy for the Northern Riffleshell, Snuffbox, Round Pigtoe, Salamander Mussel, and Rayed Bean in Canada [Proposed]. In *Species at Risk Act Recovery Strategy Series*, Fisheries and Oceans Canada, Ottawa. x + 98 p.

- Dove, C., S. Painter, and J. Kraft, 2002. Sediment quality in Canadian Lake Erie tributaries, a screening level survey. Ecosystem Health Division, Ontario Region, Environmental Conservation Branch, Environment Canada. Report No. ECB/EHD-OR/02-05/I.
- Dove, A., S. Painter, and J. Kraft. 2003. Sediment quality in Canadian Lake Ontario tributaries: A screening-level survey. Ecosystem Health Division, Ontario Region, Environmental Conservation Branch, Environment Canada, Report No. ECB/EHD-OR/03-01/I.
- Fitzsimons, J., B. Williston, G. Williston, G. Bravener, J.L. Jonas, R.M. Claramunt, J.E. Marsden, and B.J. Ellrott. 2006. Laboratory estimates of salmonine egg predation by Round Gobies (*Neogobius melanostomus*), sculpins (*Cottus cognatus* and *C. bairdi*), and crayfish (*Orconectes propinquus*). *Journal of Great Lakes Research* 32: 227–241.
- Fuller, S.L.H. 1974. Clams and mussels (Mollusca: Bivalvia). In *Pollution Ecology of Freshwater Invertebrates*. Edited by C.W. Hart, Jr. and S.L.H. Fuller. Academic Press, New York, New York, U.S.A. xiv + 389 p.
- Gagné, F., B. Bouchard, C. André, E. Farcy, and M. Fournier. 2011. Evidence of feminization in wild *Elliptio complanata* mussels in the receiving waters downstream of a municipal effluent outfall. *Comparative Biochemistry and Physiology Part C: Toxicology and Pharmacology* 153: 99–106.
- Ghedotti M.J., J.C. Smihula, and G.R. Smith. 1995. Zebra Mussel predation by Round Gobies in the laboratory. *Journal of Great Lakes Research* 21: 665–669.
- Gillis, P.L. 2011. Assessing the toxicity of sodium chloride to the glochidia of freshwater mussels: implications for salinization of surface waters. *Environmental Pollution* 159: 1702–1708.
- Gillis, P.L. 2012. Cumulative impacts of urban runoff and municipal wastewater effluents on wild freshwater mussels (*Lasmigona costata*). *Science of the Total Environment* 431: 348–356.
- Gillis, P.L., S.K. Higgins, and M.B. Jorge. 2014. Evidence of oxidative stress in wild freshwater mussels (*Lasmigona costata*) exposed to urban-derived contaminants. *Ecotoxicology and Environmental Safety* 102: 62–69.
- Gillis, P.L. and G.L. Mackie. 1994. Impact of the zebra mussel, *Dreissena polymorpha*, on populations of Unionidae (Bivalvia) in Lake St. Clair. *Canadian Journal of Zoology*, 72: 1260–1271.
- Gillis, P.L., J.C. McGeer, G.L. Mackie, M.P. Wilke, and J.D. Ackerman. 2010. The effect of natural dissolved carbon on the acute toxicity of copper to larval freshwater mussels (glochidia). *Environmental Toxicology and Chemistry* 29: 2519–2528.
- Gillis, P.L., R.J. Mitchell, A.N. Schwalb, K.A. McNichols, G.L. Mackie, C.M. Wood, and J.D. Ackerman. 2008. Sensitivity of glochidia (larvae) of freshwater mussel to copper: assessing the effect of water hardness and dissolved organic carbon on the sensitivity of endangered species. *Aquatic Toxicology* 88: 137–145.

- Hanlon, S.D. 2000. Release of juvenile mussels into a fish hatchery raceway: a comparison of techniques. Thesis (M.Sc.) Virginia Polytechnic Institute and State University, Blacksburg, Virginia.
- Holm, E., N.E. Mandrak, and M.E. Burridge. 2009. The ROM Field Guide to Freshwater Fishes of Ontario. Royal Ontario Museum, Toronto, Ontario. 462 p.
- Howells, R.G., R.W. Neck, and H.D. Murray. 1996. Freshwater Mussels of Texas. Texas Parks and Wildlife Department, Inland Fisheries Division. Austin, Texas, U.S.A. 218 p.
- Huebner, J.D. and K.S. Pynnonen. 1992. Viability of glochidia of two species of Anodonta exposed to low pH and selected metals. *Canadian Journal of Zoology* 70: 2348–2355.
- Keller, A.E. and S.G. Zam. 1990. Simplification of in vitro culture techniques for freshwater mussels. *Environmental Toxicology and Chemistry* 9: 291–1296.
- Leonard, J.A., W.G. Cope, M.C. Barnhart, and R.B. Bringolf. 2014. Metabolomic, behavioral, and reproductive effects of the synthetic estrogen 17 α -ethinylestradiol on the unionid mussel *Lampsilis fasciola*. *Aquatic Toxicology* 150: 103–116.
- Loomis, J.B. and D.S. White. 1996. Economic benefits of rare and endangered species: summary and meta-analysis. *Ecological Economics* 18: 197–206.
- Lower Great Lakes Unionid Database. 2011. Lower Great Lakes Unionid Database. Microsoft Access 2010. Department of Fisheries and Oceans, Great Lakes Laboratory of Fisheries and Aquatic Sciences, Burlington, Ontario.
- Lui, Y., W. Yang, L. Leon, I. Wong, C. McCrimmon, A. Dove, and P. Fong. 2016. Hydrologic modeling and evaluation of Best Management Practice scenarios for the Grand River watershed in Southern Ontario. *Journal of Great Lakes Research* 42(6): 1289–1301.
- Mackie, G.L. 1991. Biology of the exotic Zebra Mussel, *Dreissena polymorpha*, in relation to native bivalves and its potential impact in Lake St. Clair. *Hydrobiologia* 219: 251–268.
- Mackie, G.L. 1996. Diversity and status of Unionidae (Bivalvia) in the Grand River, a tributary of Lake Erie, and its drainage basin. Prepared for Lands and Natural Heritage Branch, Ontario Ministry of Natural Resources, Peterborough, Ontario. 39 p.
- MacDougall, T.M. and P.A. Ryan. 2012. An assessment of aquatic habitat in the southern Grand River, Ontario: water quality, lower trophic levels, and fish communities. Lake Erie Management Unit, Provincial Services Division, Fish and Wildlife Branch, Ontario Ministry of Natural Resources. Port Dover, Ontario. 141 p. + appendices.
- Metcalf-Smith, J., A. MacKenzie, I. Carmichael, and D. McGoldrick. 2005. Photo Field Guide to the Freshwater Mussels of Ontario. Published by St. Thomas Field Naturalist Club Inc. St. Thomas, ON. 60 p.

- Metcalfe-Smith, J.L., G.L. Mackie, J. Di Maio, and S.K. Staton. 2000a. Changes over time in the diversity and distribution of freshwater mussels (Unionidae) in the Grand River, southwestern Ontario. *Journal of Great Lakes Research* 26: 445–459.
- Metcalfe-Smith, J.L., D.J. McGoldrick, D.T. Zanatta, and L. Grapentine. 2007. Development of a monitoring program for tracking the recovery of endangered freshwater mussels in the Sydenham River, Ontario. WSTD Contribution, Environment Canada, Water Science and Technology Directorate, Burlington, Ontario, Canada. 40 p. + appendices.
- MOECC (Ministry of Environment and Climate Change). 1994. [Provincial Water Quality Objectives \(Ontario\)](#) (Accessed: February 2015).
- Morris, T.J. 2006. Recovery strategy for the Wavy-rayed Lampmussel (*Lampsilis fasciola*) in Canada. In *Species at Risk Act Recovery Strategy Series*. Ottawa. Fisheries and Oceans Canada. vii + 43 p.
- Morris, T.J. and A. Edwards. 2007. Freshwater mussel communities of the Thames River, Ontario: 2004–2005. Canadian Manuscript Report of Fisheries and Aquatic Science 2810. v + 30 p.
- Morris, T.J., D.J. McGoldrick, J.L. Metcalfe-Smith, D.T. Zanatta, and P.L. Gillis. 2008. Pre-COSEWIC assessment of the Wavy-rayed Lampmussel (*Lampsilis fasciola*). DFO Canadian Science Advisory Secretariat Research Document 2008/083. v + 39 p.
- Nalepa, T.F., D.J. Hartson, G.W. Gostenik, D.L. Fanslow, and G.A. Lang. 1996. Changes in the freshwater mussel community of Lake St. Clair: from Unionidae to *Dreissena polymorpha* in eight years. *Journal of Great Lakes Research* 22: 354–369.
- NatureServe. 2015. [NatureServe explorer](#): an online encyclopedia of life (web application). Version 7.1. NatureServe, Arlington, Virginia. (Accessed: February 2015).
- Neves, R.J. and M.C. Odom. 1989. Muskrat predation on endangered freshwater mussels in Virginia. *Journal of Wildlife Management* 53: 934–941.
- Nichols S.J., G. Kennedy, E. Crawford, J. Allen, J. French III, G. Black, M. Blouin, J. Hickey, S. Chernyák, R. Haas, and M. Thomas. 2003. Assessment of Lake Sturgeon (*Acipenser fulvescens*) spawning efforts in the lower St. Clair River, Michigan. *Journal of Great Lakes Research* 29: 383–391.
- Österling M.E., B.L. Arvidsson, and L.A. Greenberg. 2010. Habitat degradation and the decline of the threatened mussel *Margaritifera margaritifera*: Influence of turbidity and sedimentation on the mussel and its host. *Journal of Applied Ecology* 47: 759–768.
- Pandolfo, T.J., W.G. Cope, G.B. Young, J.W. Jones, D. Hua, and S.F. Lingenfelter. 2012. Acute effects of road salts and associated cyanide compounds on the early life stages of the unionid mussel *Villosa iris*. *Environmental Toxicology and Chemistry* 31: 1801–1806.
- Parmalee, P.W. and A.E. Bogan. 1998. *The Freshwater Mussels of Tennessee*. The University of Tennessee Press, Knoxville. 328 p.

- Poos, M., A.J. Dextrase, A.N. Schwalb, and J.D. Ackerman. 2010. Secondary invasion of the round goby into high diversity Great Lakes tributaries and species at risk hotspots: potential new concerns for endangered freshwater species. *Biological Invasions* 12: 1269–1284.
- Portt, C., G. Coker, and K. Barrett. 2007. Recovery strategy for fish species at risk in the Grand River in Canada [Draft]. In *Species at Risk Act Recovery Strategy Series*. Ottawa: Fisheries and Oceans Canada. 104 p.
- Ray, W.J. and L.D. Corkum. 1997. Predation of Zebra Mussels by Round Gobies, *Neogobius melanostomus*. *Environmental Biology of Fishes* 50: 267–273.
- Roseman E.F., W.W. Taylor, D.B. Hayes, A.L. Jones, and J.T. Francis. 2006. Predation on Walleye eggs by fish on reefs in western Lake Erie. *Journal of Great Lakes Research* 32: 415–423.
- Schloesser, D.W. and T.F. Nalepa. 1994. Dramatic decline of unionid bivalves in offshore waters of western Lake Erie after infestation by the Zebra Mussel, *Dreissena polymorpha*. *Canadian Journal of Fisheries and Aquatic Sciences* 51: 2234–2242.
- Schloesser, D.W., T.F. Napela, and G.L. Mackie. 1996. Zebra Mussel infestation of unionid bivalve (Unionidae) in North America. *American Zoologist* 36: 300–310.
- Spooner, D.E. 2007. An integrative approach to understanding the structure and function of mussel communities. Thesis (PhD) University of Oklahoma, Norman, Oklahoma.
- Stanfield, L. and R. Kuyvenhoven. 2005. Protocol for applications used in the Aquatic Landscape Inventory Software application for delineating, characterizing and classifying valley segments within the Great Lakes basin. Ontario Ministry of Natural Resources Report, July 27, 2005.
- Steinhart G.B., E.A. Marschall, and R.A. Stein. 2004. Round Goby predation on Smallmouth Bass offspring in nests during simulated catch-and-release angling. *Transactions of the American Fisheries Society* 133: 121–131.
- Strayer, D.L. and A.R. Fetterman. 1999. Changes in the distribution of freshwater mussels (Unionidae) in the upper Susquehanna River basin, 1955–1965 to 1996–1997. *American Midland Naturalist* 142: 328–339.
- Surber, T. 1913. Notes on the natural hosts of freshwater mussels. *Bulletin of the U.S. Bureau of Fisheries (Document 778)* 32: 110–116.
- Taylor, I., B. Cudmore, C. MacCrimmon, S. Madzia, and S. Hohn. 2004. The Thames River watershed: synthesis report (draft). Prepared for the Thames River Recovery Team. 74 p.
- TRRT (Thames River Recover Team). 2005. Recovery strategy for the Thames River aquatic ecosystem: 2005–2010. November 2005 Draft. 146 p.

- Tiemann, J.S., H.R. Dodd, N. Owens, and D.H. Wahl. 2007. Effects of lowhead dams on unionids in the Fox River, Illinois. *Northeastern Naturalist* 14(1): 125–138.
- Todd, A.K. and M.G. Kaltenecker. 2012. Warm season chloride concentrations in stream habitats of freshwater mussel species at risk. *Environmental Pollution* 171: 199–206.
- Tremblay, M. 2012. An effect of the invasive Round Goby (*Neogobius melanostomus*) on the recruitment of unionid mussel species at risk (Bivalvia: Unionidae). Thesis (MSc), University of Guelph. 94 p.
- USFWS (United States Fish and Wildlife Service). 1994. Clubshell (*Pleurobema clava*) and Northern Riffleshell (*Epioblasma torulosa rangiana*) recovery plan. Hadley Massachusetts. 68 p.
- UTRCA (Upper Thames River Conservation Authority). 2012. [Boaters Beware!](#). (Accessed: February 2015).
- Vaughn, C.C. and C.M. Taylor. 1999. Impoundments and the decline of freshwater mussels: a case study of an extinction gradient. *Conservation Biology* 13: 912–920.
- Vaughn, C.C. and C.C. Hakenkamp. 2001. The functional role of burrowing bivalves in freshwater ecosystems. *Freshwater Biology* 46: 1431–1446.
- Vaughn, C.C., K.B. Gido, and D.E. Spooner. 2004. Ecosystem processes performed by unionid mussels in stream mesocosms: Species roles and effects of abundance. *Hydrobiologia* 527: 35–47.
- Vaughn, C.C. and D.E. Spooner. 2006. Unionid mussels influence macroinvertebrate assemblage structure in stream. *Journal of the North American Benthological Society* 25: 691–700.
- Wang, N., C.G. Ingersoll, D.K. Hardesty, C.D. Ivey, J.L. Kunz, T.W. May, F.J. Dwyer, A.D. Roberts, T. Augspurger, C.M. Kane, R.J. Neves, and C. Barnhart. 2007. Acute toxicity of copper, ammonia, and chlorine to glochidia and juveniles of freshwater mussels (Unionidae). *Environmental Toxicology and Chemistry* 26: 2036–2047.
- Wang, N., C.G. Ingersoll, J.L. Kunz, W.G. Brumbaugh, C.M. Kane, R.B. Evans, S. Alexander, C. Walker, and S. Bakaletz. 2013. Toxicity of sediments potentially contaminated by coal mining and natural gas extraction to unionid mussels and commonly tested benthic invertebrates. *Environmental Toxicology and Chemistry* 32: 207–221.
- Watters, G.T. 2000. Freshwater mussels and water quality: a review of the effects of hydrologic and instream habitat alterations. *First Freshwater Mollusk Conservation Society Symposium*. *Ohio Biological Survey* 261–274.
- Watters, G.T., S.H. O'Dee, and S. Chordas III. 2001. Patterns of vertical migration in freshwater mussels (Bivalvia: Unionidae). *Journal of Freshwater Ecology* 16(4): 541–549.

Watters, G.T., M.A. Hoggarth, and D.H. Stansbery. 2009. *The Freshwater Mussels of Ohio*. Ohio State University Press, Columbus, Ohio. xiii + 421 p.

Wilson, C.B. 1916. Copepod parasites of freshwater fishes and their economic relations to mussel glochidia. *Bulletin of the U.S. Bureau of Fisheries* 34: 333–374.

WQB (Water Quality Branch). 1989a. The application of an interdisciplinary approach to the selection of potential water quality sampling sites in the Grand River basin. Environment Canada, Water Quality Branch, Ontario Region. 111 p.

WQB (Water Quality Branch). 1989b. The application of an interdisciplinary approach to the selection of potential water quality sampling sites in the Thames River basin. Environment Canada, Water Quality Branch, Ontario Region: 122 p.

Appendix A: effects on the environment and other species

In accordance with the [Cabinet Directive on the Environmental Assessment of Policy, Plan and Program Proposals](#) (2010), *Species at Risk Act* (SARA) recovery planning documents incorporate Strategic Environmental Assessment (SEA) considerations throughout the document. The purpose of a SEA is to incorporate environmental considerations into the development of public policies, plans, and program proposals to support environmentally sound decision-making and to evaluate whether the outcomes of a recovery planning document could affect any component of the environment or achieve any of the [Federal Sustainable Development Strategy](#)'s goals and targets.

Recovery planning is intended to benefit species at risk and biodiversity in general. However, it is recognized that strategies may also inadvertently lead to environmental effects beyond the intended benefits. The planning process based on national guidelines directly incorporates consideration of all environmental effects, with a particular focus on possible impacts upon non-target species or habitats. The results of the SEA are incorporated directly into the strategy itself, but are also summarized below in this statement.

This combined recovery strategy and action plan will clearly benefit the environment by promoting the recovery of the Fawnsfoot and Threehorn Wartyback. In particular, it will encourage the protection and improvement of riverine habitats. These habitats support species at risk from many other taxa (including birds, reptiles, fishes and plants) and thus the implementation of recovery actions for the Fawnsfoot and Threehorn Wartyback will contribute to the preservation of biodiversity in general. The potential for these recovery actions to inadvertently lead to adverse effects on other species was considered. The SEA concluded that the implementation of this document will clearly benefit the environment and will not entail any significant adverse environmental effects.

Appendix B: record of cooperation and consultation

Recovery strategies and action plans are to be prepared in cooperation and consultation with other jurisdictions, organizations, affected parties and others as outlined in *Species at Risk Act* (SARA) sections 39 and 48. Fisheries and Oceans Canada (DFO) has utilized a process of species expert/subject matter expert review to seek input to the development of this recovery strategy and action plan. Information on participation is included below.

Subject matter expert reviewers

Member / attendee	Affiliation
Todd Morris	DFO Science
Kelly McNichols-O'Rourke	DFO Science
Dave Balint	DFO Species at Risk Program
Scott Gibson	Ontario Ministry of Natural Resources
Scott Reid	Ontario Ministry of Natural Resources
Sarah Parna	Ministry of the Environment, Parks and Conservation
Daryl McGoldrick	Environment and Climate Change Canada
Patricia Gillis	Environment and Climate Change Canada
Gerry Mackie	University of Guelph
Dr. Frederick W. Schueler	Bishop Mills Natural History Centre
Erin Carroll	St. Clair Region Conservation Authority

Additional stakeholder, Indigenous, and public input will be sought through the publication of the proposed document on the Species at Risk Public Registry for a 60-day public comment period. Comments received will inform the final document.